

6th Edition, 2024

Guam Fruit and Vegetable Pesticide Guide



UNIVERSITY OF GUAM
COOPERATIVE EXTENSION
& OUTREACH

Preface

The intention of this pesticide guide is to assist farmers in selecting the proper pesticides for individual problems. The guide will be updated as needed due to the constant revisions in labeling being made by chemical manufacturers and US EPA. The guide will also be revised as new pests/pesticides are introduced into Guam.

This pesticide guide is only intended to supplement pesticide labels! Growers should always read the label first before using any pesticide.

Since the majority of pesticide registration is conducted by chemical manufacturers in the mainland United States, certain insect pests that attack plants on Guam are not found on product labels. To aid growers with insect pests not found on product labels, some suggested uses of pesticides are listed in this publication. According to federal law, it is permissible to use a particular pesticide as long as the pesticide is registered for use on that particular crop, the recommended label rate is not exceeded and spraying intervals and days to harvest are followed.

Most pesticides rates are listed in terms of pounds per acre (lbs./A) or pints per acre (pts/A). For many pesticide applicators this measurement is confusing. When preparing a 50-100 gallon tank of pesticide, please refer to Appendix A (Calibration) to determine how much pesticide should be mixed with a particular quantity of water. When mixing small amounts of pesticide, such as 1-3 gallons, please refer to Appendix B (Converting pints/acre to teaspoons/gallon). The appendices are located at the back of this guide. When mixing small amounts of powder formulations, it is very important to use an accurate weighing apparatus, such as an electronic scales or balance, in order to get the correct amount. Powders vary in concentration depending on the amount of the active ingredient as well as inert ingredients that make up the formulation.

We have also included a section (Appendix E) that provides information on Private and Commercial Certification and Recertification for the application of restricted use pesticides.

Table of Contents

Table of Contents	3
Appendix A: Calibration	5
Appendix B: Conversion and Dilution for Pesticides	7
Appendix C: Useful Conversion Factor	9
Appendix D: Core (Pesticide Core Examination), Private and Commercial Pesticide Applicator Certification for Restricted Use Pesticide (RUP) and Recertification Guidelines	11
ALIAS 4F	13
AMDRO FIRE ANT BAIT GRANUALS	18
BRAVO 720	19
CROP OIL CONCENTRATE	25
DIBROM 8 EMULSIVE	26
DIMETHOATE 4 EC	29
DIPEL DF **	32
D.Z.N. DIAZINON AG500	36
DEADLINE GT	44
EQUIS 720	46
KOCIDE 3000 **	55
LANNATE LV	63
MALATHION 5 EC	72
MALATHION 8 AQUAMUL	78
MANZATE PRO STICK	84
NEEM OIL **	90
PERM-UP 3.2 EC	91
PURESPRAY GREEN OIL **	97
SAFER INSECTICIDAL SOAP **	98
SEVIN 4F	102
SEVIN SL	111
Sluggo SNAIL & SLUG BAIT **	115
TRIACT 70 **	117
VENDEX 50WP	118
VYDATE L	119
Crop and Pest Use Index	124

TABLE OF CONTENTS

Avocado	124
Banana	126
Beans	127
Betel Nut	129
Bitter Melon	130
Broccoli	131
Cabbage	133
Cacao	135
Cantaloupe	136
Carrot	138
Cashew	140
Cauliflower	141
Chinese Cabbage	143
Corn, Field	145
Cucumber	147
Eggplant	149
Garlic	151
Gourds	153
Grapefruit	154
Green Onion	156
Guava	157
Honeydew	158
Horseradish	160
Iba	161
Jackfruit	162
Kondot	163
Lemon	164
Lettuce, Head	166
Lettuce, Leaf	168
Lime	170
Mango	172
Muskmelon	173
Mustard Greens	175
Okra	177
Orange	179
Papaya	181
Parsley	182
Peanut	184
Pepino	185
Pepper, Bell	187
Pepper, Hot	189
Pineapple	191
Pomegranate	192
Pumpkin	193
Radish	195
Squash, Summer	196
Squash, Winter	199
Sweet Potato	201
Tangerine	202
Taro	204
Tomato	205
Watermelon	207
Yam	209

OMRI Listed® ** - The Organic Materials Review Institute (OMRI) or OMRI Listed® seal assures the suitability of products for certified organic production, handling, and processing.

Read label for restrictions

Appendix A: Calibration

How to Calibrate a Hand Sprayer:

1. Measure and mark a small test area 18.5 feet X 18.5 feet. (342.25 square feet) is 1/128 of an acre.
2. Determine the output of the hand sprayer by filling a known volume of plain water to the full mark of an empty hand sprayer.
3. After pressurizing the hand sprayer, determine how long it takes in seconds to spray the marked off area with water used in # 1.
4. Fill the sprayer up again with the known volume of plain water used in # 2.
5. At the same pressure used in #3, collect the spray from the nozzle using fluid ounces (fl. oz.) in a container, for the same length of time (seconds) needed to spray the measured area. The fluid ounces collected in the container will be equal to the gallons of spray you would have applied had you sprayed a full acre (GPA).

Determine the Amount of Chemical to Add to the Tank

6. Divide the useful capacity of the tank (step 2) by the spray output, GPA (step 5) to determine the decimal fraction of an acre covered by each tank of spray.
7. Read the label to distinguish if the chemical is a concentrate or a spray solution and determine the volume of product recommended per acre. (Chemical concentrates contain large amounts of an active ingredient and an agent that requires it to be diluted with water. A spray solution is a mixture of substances that has already been diluted with water.)
8. Multiply the volume of product per acre (step 7) by the fractional acre covered per tank (step 6) to determine the amount of product to add per tank of spray.

Example:

- John measured the volume of the spray tank and found it to be two gallons. (Step 1)
- He measured and marked a test area 18.5 x 18.5 feet. (Step 2)
- He sprayed the test area and found it took 32 seconds. (Step 3)
- He caught the output in a container for 32 seconds and measured 34 fluid ounces of water caught. This shows John was putting out the equivalent of 34 gallons per acre. (Step 4)
- Each tank will cover $2 \text{ gallons} / 34 \text{ gallons} / \text{acre} = 0.059 \text{ acre}$ (Step 5)
- The product label recommends 1 quart (32 fl.-oz.) of herbicide per acre. (Step 6)

APPENDIX A

- The amount of product to add to a full spray tank is: 32 fl.-oz. per acre x 0.059 acres per tank = 1.9 fl.-oz. per spray tank. (Step 7)

This methodology works because the test area (18.5 feet squared) is 1/128 of an acre. A U.S. gallon is 128 fluid ounces. The fluid ounces of spray required to treat the test area therefore is equivalent to the gallons of spray output that would be applied to a full acre, provided the operator maintains a consistent pattern of spray output and spray coverage.

You'll need to check from time to time to make sure your sprayer is delivering the same amount of spray you calibrated. If your sprayer is delivering too much, excessive residue can accumulate on plants and you can run the risk of burning your plants with chemical sprays. If your sprayer is not delivering enough, you probably won't kill the pests you're trying to control.

Changing Delivery Rate:

There are 3 ways to change the delivery rate of your sprayer; the following methods can be used:

1. Change the pump pressure. (Lower pressure means less spray is being delivered)
2. Change the speed (e.g., tractor speed or how fast you walk) of your sprayer. Slower speed means more spray is being delivered.
3. Change the nozzles on the sprayer. The larger the hole in the nozzle, the greater the amount of spray being delivered.

Citation: *Calibrating hand-held Lancaster County Sprayers*. (n.d.). Retrieved October 27, 2022, from <https://lancaster.unl.edu/ag/factsheets/handspray.pdf>

Appendix B: Conversion and Dilution for Pesticides

CONVERTING PINTS/ACRE TO TEASPOONS/GALLON

BASED ON 100 GALLONS OF WATER/ACRE

RECOMMENDED RATE per acre (A) for LIQUIDS	CONVERTS TO (amount/gallon)
5 2/6 fluid ounces (fl oz)	1/4 teaspoons (t)
1/2 pint (pt)	8 fluid ounces (fl oz)
10 2/6 fluid ounces (fl oz)	2/3 teaspoons (t)
1 pint (pt)	1 teaspoon (t)
1 1/2 pints (pt)	1 1/2 teaspoons (t)
1 quarts (qt)	2 teaspoons (t)
2 1/2 pints (pt)	2 1/2 teaspoons (t)
3 pints (pt)	1 Tablespoon = 3 teaspoons
1 2/3 quarts (qt)	3 1/4 teaspoons (t)
3 1/2 pints (pt)	3 1/2 teaspoons (t)
4 pints (pt)	4 teaspoons (t)
4 1/2 pints (pt)	4 1/2 teaspoons (t)
5 pints (pt)	5 teaspoons (t)
5 1/2 pints (pt)	5 1/2 teaspoons (t)
6 pints (pt)	2 Tablespoons = 6 teaspoons
4 quarts (qt)	8 teaspoons (t)
8 1/4 pints (pt)	8 1/4 teaspoons (t)

APPENDIX B

DILUTION CHART FOR PREPARING LESS THAN 100 GALLONS OF PESTICIDE SPRAY

Formulation	100 gal	3 gal	1 gal
Emulsifiable Concentrate	1 pt	1 tsp	1 tsp
	1 qt	2 tbsp	2 tsp
	1 gal	8 tbsp	3 tbsp
	2 gal	1/2 pt	5 tbsp
	3 gal	3/4 pt	1/4 pt
	4 gal	1 pt	10 tbsp
	5 gal	1 1/4 pt	13 tbsp
Wettable Powder	1/2 lb	1 tsp	1 tsp
	1 lb	6 tsp	2 tsp
	2 lbs	4 tbsp	4 tsp
	3 lbs	6 tbsp	2 tbsp
	4 lbs	8 tbsp	8 tsp
	5 lbs	9 tbsp	3 tbsp

Appendix C: Useful Conversion Factor

Liquid		
1 Tablespoon (T)	=	3 teaspoons (t)
1 fluid ounce (fl oz)	=	2 Tablespoons (T)
1 cup (C)	=	8 ounces (fl oz)
1 pint (pt)	=	16 fluid ounces (fl oz)
1 quart (qt)	=	4 cups (C)
4 cups (C)	=	2 pints (pt)
4 cups (C)	=	32 fluid ounces (fl oz)
2 pints (pt)	=	32 fluid ounces (fl oz)
1 gallon (gal)	=	3.785 liters (L)
1 gallon (gal)	=	128 fluid ounces (fl oz)
3.785 liters (L)	=	128 fluid ounces (fl oz)
1 quart (qt)	=	946 milliliters (ml)
1 fluid ounce (fl oz)	=	29.54 milliliters (ml)
1 liter (L)	=	1.06 quarts (qt)
1 liter (L)	=	0.264 gallons (gal)
Dry		
1 ounce (oz)	=	28.35 grams (g)
1 pound (lb)	=	453.59 grams (g)
1 pound (lb)	=	16 ounces (oz)
453.59 grams (g)	=	16 ounces (oz)
1 ton (t)	=	2000 pounds (lb)
1 hundredweight	=	100 pounds (lb)
1 gram (g)	=	0.0353 oz
1 kilogram (kg)	=	2.205 (lb)
1 kilogram (kg)	=	1000 grams (g)
1 metric ton (MT)	=	1.1 tons (t)

APPENDIX C

Length		
1 inch (in)	=	2.54 centimeters (cm)
1 foot (ft)	=	0.35 meters (m)
1 yard (yd)	=	3 feet (ft)
1 centimeter (cm)	=	0.394 inches (in)
1 meter (m)	=	3.28 feet (ft)
Area		
1 acre (A)	=	43,560 square feet (sq ft)
1 acre (A)	=	0.405 hectare (ha)
1 hectare (ha)	=	2.47 acres (A)
Yield or Rate		
1 lb/A	=	1.12 kg/ha
1 kg/ha	=	0.89 lb/A
Temperature		
°F	=	$\frac{9}{5} \text{ } ^\circ\text{C} + 32$
°C	=	$\frac{5}{9} (\text{ } ^\circ\text{F} - 32)$

Appendix D: Core (Pesticide Core Examination), Private and Commercial Pesticide Applicator Certification for Restricted Use Pesticide (RUP) and Recertification Guidelines

CORE

All individuals seeking Private or Commercial Pesticide Applicator certification must pass the PESTICIDE CORE examination. A pesticide study manual entitled “Applying Pesticides Correctly: A Guide for Private and Commercial Applicators” (CORE GUIDE) contains all the information you will need to read and understand before you can successfully pass the PESTICIDE CORE examination. This pesticide manual contains 11 units. Each unit begins with learning objectives, terms to know, and at the end of each unit a set of questions that you need to understand and eventually be able to answer on your own. Also, P. L. 30-197 states that Core Training is required for any applicant who intends to use general use pesticides as part of carrying out his occupational duties. The Core Training consists of sixteen (16) hours of training.

The information contained in this pesticide manual provides both the Private and Commercial Applicator the basic knowledge needed to pass the PESTICIDE CORE examination. A score of 70% or above is considered a passing score. Upon request, the PESTICIDE CORE examination can be taken at the Guam Environmental Protection Agency (GEPA). Once you successfully pass the PESTICIDE CORE examination, two (2) options are available to you:

Option 1. Private Applicator for (RUP)

Any individual that has successfully passed the PESTICIDE CORE examination with a score of 70% or above and completes the sixteen (16) hour training course can participate in a Private Pesticide workshop. P. L. 30-197 states that Category Training is required for all restricted use pesticide applicators. Satisfactory completion of Core Training is required as a prerequisite for Category Training. Category Training consists of sixteen hours (16) of a specialized type of training course and a closed-book exam. A pesticide study manual entitled “Applying Pesticides Correctly: A Supplement Guide for Private Applicators” (PRIVATE GUIDE) is available for this category. GEPA defines Private Applicators as “persons using or supervising restricted use pesticides for the purpose of producing any agricultural commodity on property owned or rented by him or his employer, or, if applied without compensation other than trading of personal services between producers of agricultural commodities on the property of another person. Upon request, the PRIVATE PESTICIDE examination can be taken at the Guam Environmental Protection Agency (GEPA).

Option 2. Commercial Applicator for (RUP)

Any individual that has successfully passed the PESTICIDE CORE examination with a score of 70% or above and completes the sixteen (16) hour training course can participate in a Commercial Pesticide workshop. P. L. 30-197 states that Category Training is required for all restricted use pesticide applicators. Satisfactory completion of Core Training is required as a prerequisite for Category Training. Category Training consists of sixteen hours (16) of a specialized type of training course and a closed-book exam. Guam EPA defines Commercial Applicators as “persons applying or supervising the use of restricted use pesticides in pest control work, demonstration work or similar pest control activity”. Upon request, the Commercial applicator examination can be taken at the Guam Environmental Protection Agency (GEPA).

(RUP) - Applicator Categories:

1. Agriculture Pest Control
 - 1.1.Plants
 - 1.2.Animals
2. Forest Pest Control
3. Ornamental and Turf Pest Control
4. Seed Treatment
5. Aquatic Pest Control
6. Right-of-way Pest Control
7. Industrial, Institutional, Structural and Health-Related Pest Control
 - 7.1.General and Structural Pest Control
 - 7.2.Structural and Commodity Fumigation
 - 7.3.Soil Fumigation
 - 7.4.Chlorine Gas
8. Public Health Pest Control
9. Regulatory Pest Control

RENEWAL of Core, Private, and Commercial Pesticide Applicator Cards

All Certification Cards shall be renewed every three (3) years. Individuals may submit applications for renewal no more than thirty (30) days prior to the expiration date of their certification. Individuals intending to renew their certification must successfully pass the appropriate exam with a score of 70% or above.

Any questions concerning Guam’s Pesticide laws please call the Guam Environmental Protection Agency at 671-300-4751/2/3.

ALIAS 4F

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Banana	Aphids Leafhoppers Thrips	3.2 oz/A	0 days	14 days
<p>Specific Directions: *Maximum Alias 4F allowed per crop season when making foliar applications: 16 fluid ounces per acre *Apply as a broadcast or directed foliar spray to area as pest populations build *Two applications may be required *Read the label</p>				
Brassica Leafy Vegetables: Broccoli Cabbage Cauliflower Chinese Cabbage Mustard Greens	Aphids Flea Beetles Leafhoppers Whiteflies	1.5 oz/A	7 days	5 days
<p>Specific Directions: *Maximum Alias 4F allowed per crop season when making foliar applications: 7.68 fluid ounces per acre *Apply as a broadcast or directed foliar spray to area as pest populations build *Two applications may be required *Not for use on crops grown for seed unless allowed by state-specific labeling</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Citrus: Grapefruit Lemon Lime Tangerine Orange	Aphids Black Flies Flea Beetles Leafhoppers Leafminers Mealybugs Scales Sharpshooters Thrips (Suppressed) Whiteflies	4-8 oz/A	0	10 days
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Maximum Alias 4F allowed per crop season when making foliar applications: 16 fluid ounces per acre *Apply as a broadcast or directed foliar spray to area as pest populations build *Two applications may be required *Do not apply during bloom or within 10 days prior to bloom or when bees are foraging 				
Cucurbits: (SOIL TREATMENT) Bitter melon Cantaloupe Gourd Honeydew Muskmelon Pumpkin Squash (Summer, Winter) Watermelon	Aphids Cucumber Beetles Leafhoppers Thrips Whiteflies	8-12 oz/A	21	See label
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Maximum Alias 4F allowed per crop season when making foliar applications: 12 fluid ounces per acre *Not for use on crops grown for seed unless allowed by state-specific labeling *Read the label for application methods 				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Fruiting Vegetables: Eggplant Okra Pepper (Bell, Chili, Sweet) Tomato Pepino	Aphids Colorado Potato Beetle Leafhoppers Whiteflies	1.5-2.4 oz/A	0 days	5 days
<p>Specific Directions: *Maximum Alias 4F allowed per crop season when making foliar applications: 7.68 fluid ounces per acre *Not for use on crops grown for seed unless allowed by state-specific labeling *Apply as a broadcast or directed foliar spray to area as pest populations build *Two applications may be required *Not for use on crops grown for seed unless allowed by state-specific labeling</p>				
Leafy Vegetables: Lettuce (Head, Leaf) Parsley	Aphids Flea Beetles Leafhoppers Whiteflies	1.5 oz/A	7 days	5 days
<p>Specific Directions: *Maximum Alias 4F allowed per crop season when making foliar applications: 7.6 fluid ounces per acre *Not for use on crops grown for seed unless allowed by the state *Apply as a broadcast or directed foliar spray to area as pest populations build *Two applications may be required</p>				
Legume Vegetables: Beans	Aphids Leafhoppers Whiteflies	1.4 oz/A	7 days	See label
<p>Specific Directions: *Maximum Alias 4F allowed per crop season when making foliar applications: 4.2 fluid ounces per acre *Not for use on crops grown for seed unless allowed by the state *Apply as a broadcast or directed foliar spray to area as pest populations build *Two applications may be required</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Peanut (SOIL TREATMENT)	Aphids Leafhoppers Thrips (Suppressed) Whiteflies	8-12 oz/A	14 days	See label
<p>Specific Directions: *Maximum Alias 4F allowed per crop season when making soil applications: 12 fluid ounces per acre *Prior to applying Alias 4F to peanut, consult with the State, Cooperative Extension Service, or a AFAMA representative for recommendations *Read the label for application methods</p>				
Root Vegetables: Carrot Parsley (Turnip- Rooted) Radish	Aphids Flea Beetles Leafhoppers Whiteflies	1.4 oz/A	7 days	5 days
<p>Specific Directions: *Maximum Alias 4F allowed per crop season when making foliar applications: 1.4 fluid ounces per acre on radish, 4.2 fluid ounces on other crops *Maximum Alias 4F application(s) per crop season: 1 on radish, 3 on all other crops *Not for use on crops grown for seed unless allowed by the state *Apply as a broadcast or directed foliar spray to area as pest populations build *Two applications may be required</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Tropical Fruit: Avocado Guava Mango Papaya	Aphids Leafhoppers Mealybugs Scales (Suppressed) Thrips Whiteflies	3.2 oz/A	7 days	10 days
<p>Specific Directions: *Maximum Alias 4F allowed per crop season when making foliar applications: 16 fluid ounces per acre *Do not apply pre-bloom or during bloom or when bees are foraging *Apply as a broadcast or directed foliar spray to area as pest populations build *Two applications may be required</p>				
Tuberous and Corn Vegetables: Sweet Potato Yam	Aphids Cucumber Beetles Leafhoppers Whiteflies	1.4 oz/A	7 days	5 days
<p>Specific Directions: *Maximum Alias 4F allowed per crop season when making foliar applications: 1.4 fluid ounces per acre for radish, 4.2 fluid ounces on all other crops *Not for use on crops grown for seed unless allowed by state-specific labeling *Read the label for application methods</p>				

AMDRO FIRE ANT BAIT GRANUALS

CROP	PEST(S) CONTROLLED	
<p>Tropical Fruits and Nuts:</p> <p>Avocado Banana Breadfruit Cashew Guava Mango Papaya Star Apple</p>	<p>Bighead ants Fire ants Little fire ants</p>	<p>See label</p> <p>Special Directions:</p> <ul style="list-style-type: none"> *Amdro may be applied in Tropical Fruits and nut orchards when applied in bait stations *Do not apply Amaro directly to the ground around these crops *Do not apply directly to any part of the plant *When applying for control of little fire ants, apply Amaro in orchards only during non-fruiting periods *Remove bait stations from orchard after flowering but before fruiting structures appear on crop plants *Do not apply Amaro in orchards when fruits are presents on crop plants *Read label for bait station directions

BRAVO 720

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Beans, Dry	Anthracnose Cercospora Leaf Spot	1.375-2 pts/A	14	7-10 days
	Specific Directions: *Use in sufficient water to obtain adequate coverage *Begin applications during Early bloom stage *For use only on beans harvested dry with pods removed *Do not apply more than 4 times per growing season *Do not allow livestock to graze in treated areas or feed treated plant parts to livestock			
Beans, Snap	Botrytis Blight (Gray Mold)	3 pts/A	7 days	7 days
	Specific Directions: *Use in sufficient water to obtain adequate coverage *Begin applications during Early bloom stage or when disease first threatens *Do not graze treated areas or feed treated plant parts to livestock			
	Rust	1.375-3 pts/A	7 days	7 days
	Specific Directions: *Use in sufficient water to obtain adequate coverage *Begin applications during Early bloom stage or when disease first threatens *Do not graze treated areas or feed treated plant parts to livestock			
Broccoli Cabbage Cauliflower Chinese Cabbage	Alternaria Leaf Spot	1.5 pts/A	7 days	7-10 days
	Specific Directions: *Use in sufficient water to obtain adequate coverage *Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development			

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Carrot	Cercospora (Early) Blight Alternaria (Late) Blight	1.5-2 pts/A	0	7-10 days
	Specific Directions: *Read the label *Use in sufficient water to obtain adequate coverage *Start applications when disease threatens *Bravo 720 may be applied through sprinkler irrigation equipment			
Corn, Sweet	Helminthosporium Leaf Blight Rust	0.75-2 pts/A	14 days	4-7 days
	Specific Directions: *Use in sufficient water to obtain adequate coverage *Begin applications when conditions favor disease development *Under severe disease conditions, use 1.5-2 pts/A *Do not apply to sweet corn to be processed *Do not allow livestock to graze in treated fields *Do not ensile treated corn or use as livestock forage			
Cucurbits: Cantaloupe Cucumber Honeydew Muskmelon Pumpkin Squash, Summer Squash, Winter Watermelon	Anthracnose Target Spot	1.5-2 pts/A	0 days	7 days
	Specific Directions: *Use in sufficient water to obtain adequate coverage *Begin applications when plants are in true leaf stage or when conditions are favorable for disease development *Under severe disease conditions, shorten spray interval *Bravo 720 may be applied through sprinkler irrigation equipment *Watermelon use: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. Do not apply Bravo 720 to watermelons when any of the following conditions are present: (1) intense heat and sunlight, (2) drought conditions, (3) poor vine canopy, (4) other crop and environmental conditions which may be conducive to increased natural sunburn *Do not combine Bravo 720 with anything excepts water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use			

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Cercospora Leaf Spot Gummy Stem Blight Alternaria Leaf Blight Powdery Mildew Scab (No Scab on Cucumber)	2-3 pts/A	0 days	7 days
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water to obtain adequate coverage *Begin applications when plants are in true leaf stage or when conditions are favorable for disease development *Under severe disease conditions, shorten spray interval *Bravo 720 may be applied through sprinkler irrigation equipment *Watermelon use: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. Do not apply Bravo 720 to watermelons when any of the following conditions are present: (1) intense heat and sunlight, (2) drought conditions, (3) poor vine canopy, (4) other crop and environmental conditions which may be conducive to increased natural sunburn *Do not combine Bravo 720 with anything excepts water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use 				
Garlic Green Onion	Botrytis Blight (Gray Mold)	1.5-3 pts/A	7 (garlic) 14 (green onion)	7-10 days
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water to obtain thorough coverage of tops. *Begin applications prior to favorable infection periods, and repeat for as long as conditions favor disease *Use the high rate and a 7 day schedule of applications when heavy dew or rain persist *Do not apply more than 3 times per season in green onions. *If additional disease control is needed before harvest, use another registered fungicide 				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Papaya	Anthracnose Stem End Rot	1.5-3 pts/A	0 days	14 days
	Specific Directions: *Use in sufficient water to obtain adequate coverage of fruit and leaves *Begin treatment when conditions favor development of disease and continue treatments until weather conditions no longer favor disease development *Do not graze livestock in treated area or feed processing by-products to livestock			
Peanut	Early Leaf Spot (Cercospora) Late Leaf Spot (Cercosporidium)	1-1.5 pts/A	14 days	10-14 days
	Specific Directions: *Read the label *Apply in sufficient water for adequate coverage when leaf wetness first occurs or 30-40 days after planting *When conditions favor Late Leaf Spot or when rust occur, apply 1.5 pts/A at 10 day intervals for the remainder of the season *Do not allow livestock to graze in treated areas *Do not feed hay or threshings from treated fields to livestock *Bravo 720 may be applied through sprinkler irrigation equipment			
	Rust	1.5 pts/A	14 days	10-14 days
Specific Directions: *Read the label *Apply in sufficient water for adequate coverage when leaf wetness first occurs or 30-40 days after planting *When conditions favor Late Leaf Spot or when rust occur, apply 1.5 pts/A at 10 day intervals for the remainder of the season *Do not allow livestock to graze in treated areas *Do not feed hay or threshings from treated fields to livestock *Bravo 720 may be applied through sprinkler irrigation equipment				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Tomato	Foliage: Early Blight Botrytis Blight (Gray Mold) Gray Leaf Spot Septoria Leaf Spot Target Spot	2-2.75 pts/A	0 days	7-10 days
Specific Directions: *Read the label *Apply in sufficient water to obtain adequate coverage *Begin applications when dew or rain occur and disease threatens *Use the highest rate and shortest interval specified when disease conditions are severe *Bravo 720 may be combined in the spray tank with EPA-registered pesticide products that claim copper as the active ingredient and are labeled for control of bacterial diseases of tomatoes. Check the copper manufacturer's label for specific instructions, precautions, and limitations prior to mixing with Bravo 500 *Bravo 720 may be applied through sprinkler irrigation equipment				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Tomato	Fruit: Anthracnose Alternaria Fruit Rot (Black Mold) Botrytis Blight (Gray Mold) Rhizoctonia Fruit Rot	2-2.75 pts/A	0 days	7-14 days (beginning at fruit set)
Specific Directions: *Read the label *Apply in sufficient water to obtain adequate coverage *Begin applications when dew or rain occur and disease threatens *Use the highest rate and shortest interval specified when disease conditions are severe *Bravo 720 may be combined in the spray tank with EPA-registered pesticide products that claim copper as the active ingredient and are labeled for control of bacterial diseases of tomatoes. Check the copper manufacturer's label for specific instructions, precautions, and limitations prior to mixing with Bravo 500 *Bravo 720 may be applied through sprinkler irrigation equipment				

CROP OIL CONCENTRATE

Consult your agricultural extension agent for information on pesticide use for this crop.

DIBROM 8 EMULSIVE

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Beans	Aphids Leafhoppers Loopers Leafminers Spider Mites	1 pts/A	1 day	7 days
	Specific Directions: *Read the label *Begin application at first sign of insects *Do not feed treated vines or hay *Do not make more than 5 applications per season			
	Large Loopers Stink Bugs	1.5 pts/A	1 day	7 days
Specific Directions: *Read the label *Begin application at first sign of insects *Do not feed treated vines or hay *Do not make more than 5 applications per season				
Broccoli Cabbage Chinese Cabbage Cauliflower	Aphids Cabbageworm Diamondback Caterpillars	1 pt/A	1 day	7 days
	Specific Directions: *Read the label *Do not apply when temperature is over 90°F			
Cantaloupe Muskmelon	Aphids Armyworms Loopers Leafhoppers Leafminers	1 pts/A (minimum of 30 gals. of water)	1 day	7 days
	Specific Directions: *Read the label *Do not apply when temperature is over 90°F *Do not apply more than 2 pts of product per acre per season			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Citrus: Grapefruit Lemon Orange Tangerine	Aphids Mites Leafrollers Scale Crawlers Thrips	1-2 pts/A use 100-500 gals. of water	7 days	An Needed
Specific Directions: *Read the label *Apply when pests first appear				
Eggplant Pepper (Bell, Hot)	Aphids Beetles Leafminers Flea Beetles Spider Mites	1-2 pts/A (minimum of 20 gals. of water)	1 day	7 days
Specific Directions: *Read the label *Do not apply when temperature is over 90°F *Do not apply more than 6 pts of product per acre per season *Do not make more than 5 applications per season				
Summer Squash	Aphids Armyworms Cucumber Beetles Spider Mites Loopers Thrips	1-2 pts/A (minimum of 20 gals. of water)	1 day	7 days
Specific Directions: *Read the label *Do not apply when temperature is over 90°F *Do not apply more than 6 pts of product per acre per season				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Summer Squash	Leafminers	1.5-2 pts/A (minimum of 20 gals. of water)	1 day	7 days
	Specific Directions: *Read the label *Do not apply when temperature is over 90°F *Do not apply more than 6 pts of product per acre per season *Do not make more than 5 applications per season			

DIMETHOATE 4 EC

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Beans (Green, Snap, Dry)	Aphids Grasshoppers Beetles Leafhoppers Leafminers Mites Plant Bugs	0.5-1 pts/A	7 days	As needed
	Specific Directions: *Do not feed treated vines *This pesticide is highly toxic to bees, do not apply if bees are visiting the areas to be treated when crop or weeds are in bloom			
Broccoli Cauliflower	Aphids	0.5-1 pts/A	7 days	As needed
	Specific Directions: *Read the label			
Cabbage	Aphids	0.5-1 pts/A	3 days	As needed
	Specific Directions: *Read the label			
Citrus: Grapefruit Lemon Orange Tangerine	Aphids	0.5-1 pts per 100 gals. water	15 days	As needed
	Specific Directions: *Apply as an outside coverage spray			
	Mites	0.5-1 pts per 100 gals. water	15 days	An Needed
	Specific Directions: *Apply as an outside coverage spray			
	Scales	1-1.5 pts per 100 gals. water	45 days	An Needed
Specific Directions: *Apply as a thorough coverage spray				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Citrus: Grapefruit Lemon Orange Tangerine	Thrips	0.5-1 pts per 100 gals. water	15 days	An Needed
	Specific Directions: *Apply as a mist spray			
	Whiteflies	1 pts per 100 gals. water	15 days	An Needed
	Specific Directions: *Apply as a thorough distribution coverage spray			
Field Corn	Aphids Mites Beetles Caterpillars Spider Mites	0.67-1 pts/A	14 days	As needed
	Specific Directions: *Make no more than 3 applications per year *Do not feed or graze within 14 days of last application *Do not apply to com during the pollen-shed period			
	Grasshoppers	1 pts/A in 20-40 gals of water	14 days	As needed
	Specific Directions: *Make no more than 3 applications per year *Do not feed or graze within 14 days of last application *Do not apply to com during the pollen-shed period			
Melons: Cantaloupe Honeydew Muskmelon Pepino	Aphids Leafhoppers Leafminers	1 pts/A	3 days	As needed
	Specific Directions: *Read the label			
Lettuce, Head	Aphids Leafhoppers Leafminers	0.5 pts/A	7 days	As needed

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
<p>Specific Directions: *Read the label</p>				
<p>Lettuce, Leaf Mustard Greens</p>	<p>Aphids Leafhoppers Leafminers</p>	<p>0.5 pts/A</p>	<p>7 days</p>	<p>As needed</p>
	<p>Specific Directions: *Read the label</p>			
<p>Pepper</p>	<p>Aphids Leafminers</p>	<p>0.5-0.67 pts/A</p>	<p>0 days</p>	<p>As needed</p>
	<p>Specific Directions: *Read the label</p>			
<p>Tomato</p>	<p>Aphids Leafhoppers Leafminers</p>	<p>0.5-1 pts/A</p>	<p>7 days</p>	<p>As needed</p>
	<p>Specific Directions: *Where cabbage worms and cabbage Loopers are a problem, the above rates are compatible with endosulfan, malathion or parathion. Use in accordance with the manufacturer's directions for control of these insects</p>			
<p>Watermelon</p>	<p>Aphids Leafminers</p>	<p>0.5-1 pts/A</p>	<p>3 days</p>	<p>As needed</p>
	<p>Specific Directions: *Read the label</p>			

DIPEL DF **

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Avocado	Caterpillars Leafrollers Loopers	0.5-2 lb/A	0 days	As needed
	Special Directions: *Read the label			
Banana	Banana skipper	0.5-1 lb/A	0 days	As needed
	Special Directions: *Read the label			
Beans	Armyworms	1-2 lbs./A	0 days	As needed
	Special Directions: *DiPel DF may be used to control small Armyworms and/or podworms (1st and 2nd instar) when populations are light and full coverage sprays are applied *If mature worms or heavy populations are present a contact insecticide should be used to enhance control			
	Loopers Caterpillars	0.5-1 lb/A	0 days	As needed
	Special Directions: *Read the label			
Bitter melon Cantaloupe Cucumber Gourds Honeydew Kondot Muskmelon Pepino Pumpkin Squash, Summer Squash, Winter Watermelon	Armyworms	1-2 lbs./A	0 days	As needed
	Special Directions: *DiPel DF may be used to control small Armyworms and/or podworms (1st and 2nd instar) when populations are light and full coverage sprays are applied *If mature worms or heavy populations are present a contact insecticide should be used to enhance control			
	Loopers Melonworms	0.5-1 lb/A	0 days	An Needed
	Specific Directions: *Read the label			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Carrot	Armyworms	1-2 lbs./A	0 days	As needed
	Special Directions: *DiPel DF may be used to control small Armyworms and/or podworms (1st and 2nd instar) when populations are light and full coverage sprays are applied *If mature worms or heavy populations are present a contact insecticide should be used to enhance control			
	Caterpillars Cutworms Diamondback Moths Leafrollers Loopers	0.5-1 lb/A	0 days	As needed
	Special Directions: *Read the label			
Citrus: Grapefruit Lemon Lime Orange Tangerine	Cutworms	0.5-2 lbs./A	0 days	As needed
	Special Directions: *Apply to light to moderate populations of newly-hatched worms *Read the label			
	Leafrollers	0.5-2 lbs./A	0 days	As needed
	Special Directions: *Read the label			
Corn	Armyworms	1-2 lbs./A	0 days	As needed
	Special Directions: *DiPel DF may be used to control small Armyworms and/or podworms (1st and 2nd instar) when populations are light and full coverage sprays are applied *If mature worms or heavy populations are present a contact insecticide should be used to enhance control			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Eggplant Pepper, Bell Pepper, Hot Tomato	Armyworms	1-2 lbs./A	0 days	As needed
	Special Directions: *DiPel DF may be used to control small Armyworms and/or podworms (1st and 2nd instar) when populations are light and full coverage sprays are applied *If mature worms or heavy populations are present a contact insecticide should be used to enhance control			
	Caterpillars Cutworms Loopers	1-2 lbs./A	0 days	As needed
	Special Directions: *Read the label			
Garlic Green onion	Armyworms	1-2 lbs./A	0 days	As needed
	Special Directions: *DiPel DF may be used to control small Armyworms and/or podworms (1st and 2nd instar) when populations are light and full coverage sprays are applied *If mature worms or heavy populations are present a contact insecticide should be used to enhance control			
	Caterpillars Cutworms Diamondback Moths Leafrollers Loopers	0.5-1 lb/A	0 days	As needed
	Specific Directions: *Read the label			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Lettuce (Head, Leaf) Broccoli Cabbage Mustard Greens Cauliflower Chinese Cabbage Parsley	Armyworms	1-2 lbs./A	0 days	As needed
	Special Directions: *DiPel DF may be used to control small Armyworms and/or podworms (1st and 2nd instar) when populations are light and full coverage sprays are applied *If mature worms or heavy populations are present a contact insecticide should be used to enhance control			
	Caterpillars Diamondback Moths Cutworms Leafrollers Loopers	0.5-1 lb/A	0 days	As needed
	Specific Directions: *Read the label			
Peanut	Caterpillars Loopers	0.5-1 lb/A	0 days	As needed
	Special Directions: *Read the label			

For Smaller Spray Volume

If Rate Is	Use This Amount Per Gallon (wt)
1/4 lb./acre or 100 gals.	1/2 tsp. (0.04 oz)
1/2 lb./acre or 100 gals.	1 tsp. (0.08 oz)
1 lb./acre or 100 gals.	2 tsps. (0.16 oz)
2 lb./acre or 100 gals.	4 tsps. (0.32 oz)

D.Z.N. DIAZINON AG500

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Beans (Pole, Snap)	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			
	Aphids Beetles Leafminers	1-1.5 pts/A	7 days	5 days
	Specific Directions: *Apply as insects occur *Do not make more than 3 applications per season *Wait 4 days following treatment if forage is to be cut for hay *Diazinon will not control organophosphate-resistant Leafminers *To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom			
	Cucumber Beetles	0.75 pts/A	7 days	5 days
Broccoli	Specific Directions: *Apply as insects occur *Do not make more than 3 applications per season *Wait 4 days following treatment if forage is to be cut for hay *Make Cucumber Beetles application in 100 gals. of water per acre *To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom			
	Cutworms	2-4 qts/A	See label	See label

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Aphids Diamondback Moths Caterpillars Leafminers	0.5-1 pts/A	7 days	7 days
<p>Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season</p>				
Cabbage	Cutworms	2-4 qts/A	See label	See label
<p>Specific Directions: *Broadcast just before planting and immediately incorporate into the soil</p>				
	Aphids Diamondback Moths Caterpillars Leafminers	0.5-1 pts/A	21 days	7 days
<p>Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season</p>				
Cantaloupe Honeydew Muskmelon Watermelon	Cutworms	2-4 qts/A	See label	See label
<p>Specific Directions: *Broadcast just before planting and immediately incorporate into the soil</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Aphids Cucumber Beetles Leafhoppers Leafminers Melonworms Spider Mites Thrips	0.5-1.5 pts/A	3 days	7 days
	Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season			
Carrot	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			
	Aphids	1 pts/A	14 days	7 days
	Specific Directions: *Apply as insects occur *Do not make more than 5 applications per season			
Cauliflower	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			
	Aphids Diamondback Moths Caterpillars Leafminers	0.5-1 pts/A	5 days	7 days
	Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Chinese Cabbage	Aphids Diamondback Moths Caterpillars Leafminers	0.5-1 pts/A	10 days	7 days
	Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season			
Cucumber	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			
	Aphids Diamondback Moths Caterpillars Leafminers	1 pts/A	7 days	7 days
	Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season			
Lettuce (Head, Leaf)	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			
	Aphids Leafminers	0.5-1 pts/A	14 days	7 days
	Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season			
Mustard Greens	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Aphids Diamondback Moths Caterpillars Leafminers	0.5-1 pts/A	10 days	7 days
<p>Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season</p>				
Parsley	Cutworms	2-4 qts/A	See label	See label
<p>Specific Directions: *Broadcast just before planting and immediately incorporate into the soil</p>				
Pepper (Bell, Hot)	Cutworms	2-4 qts/A	See label	See label
<p>Specific Directions: *Broadcast just before planting and immediately incorporate into the soil</p>				
	Aphids Leafminers	0.5-1 pts/A	5 days	7 days
<p>Specific Directions: *Apply as insects occur. *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season</p>				
Radish	Cutworms	2-4 qts/A	See label	See label
<p>Specific Directions: *Broadcast just before planting and immediately incorporate into the soil</p>				
	Aphids Flea Beetles Leafminers	0.5-1 pts/A	14 days	7 days
<p>Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Squash, Summer	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			
	Aphids Cucumber Beetles Leafhoppers Leafminers Melonworms Spider Mites Thrips	0.5-1.5 pts/A	7 days	7 days
	Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season			
Squash, Winter	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			
	Aphids Cucumber Beetles Leafhoppers Leafminers Melonworms Spider Mites Thrips	0.5-1.5 pts/A	3 days	7 days
	Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season			
Sweet Potato	Flea Beetles	3-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Sweet Corn	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			
	Caterpillars	1-1.25 qts/A	See label	7 days
	Specific Directions: *Apply when silks first appear *Do not make more than 5 applications per season			
	Aphids	1-2 pts/A	See label	7 days
	Specific Directions: *Apply in sufficient water to provide complete coverage to the upper portions of the plants *Do not make more than 5 applications per season			
	Beetles	2-2.5 pts/A	See label	7 days
	Specific Directions: *Do not make more than 5 applications per season			
	Flea Beetles Grasshoppers Spider Mites	1 pts/A	See label	7 days
	Specific Directions: *Apply as insects occur *Do not make more than 5 applications per season			
SWEET CORN NOTES: (1) To protect bees, do not apply foliarly to corn during the pollen shed period (2) Allow a minimum of 7 days between the last application and grazing or cutting for forage (3) Corn may be picked 7 days after the last application				
Tomato	Cutworms	2-4 qts/A	See label	See label
	Specific Directions: *Broadcast just before planting and immediately incorporate into the soil			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Aphids Leafminers	0.5 pts/A	1 day	7 days
	Specific Directions: *Apply as insects occur *Diazinon will not control organophosphate-resistant Leafminers *Do not make more than 5 applications per season			
	Armyworms Cucumber Beetles	0.75-1 pts/A	1 day	7 days
	Specific Directions: *Apply as insects appear *Do not make more than 5 applications per season			

DEADLINE GT

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Citrus	Slugs Snails	25lbs./A	0 days	14 days
	Specific Directions: *Total number of applications per season: 6 *Read the label for application directions			
Corn, Field	Slugs Snails	25lbs./A	0 days	7 days
	Specific Directions: *Total number of applications per season: 3 *Read the label for application directions			
Leafy Greens: Broccoli Cabbage Cauliflower Collards Kale Mustard Greens	Slugs Snails	25lbs./A	0 days	14 days
	Specific Directions: *Total number of applications per season: 3 *Read the label for application directions			
Lettuce	Slugs Snails	25lbs./A	0 days	14 days
	Specific Directions: *Total number of applications per season: 3 *Read the label for application directions			
Tomato	Slugs Snails	25lbs./A	0 days	14 days
	Specific Directions: *Total number of applications per season: 3 *Read the label for application directions			

Restrictions

- This product is intended solely for use on agricultural crops grown for commercial or research purposes, and in State and/or Federal invasive mollusk eradication operations.
- This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other sites such as the exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, recreation areas, etc.
- Do not apply as wide-area broadcast. Broadcast applications and applications of this product directed to plant parts are prohibited except to grass grown for seed, taro and as specified for use in State and/or Federal invasive mollusk eradication operations.
- Do not allow bait pellets to come in contact with plant parts except to grass grown for seed, taro, and as specified for use in State and/or Federal invasive mollusk eradication operations.
- Use only as a baiting treatment to be applied directly to the soil area beneath listed plants.
- Read entire label before using.
- Use as band treatment between rows.
- Do not apply to dry soil. Make applications following irrigation.
- Do not re-water for 48 hours after application.
- Apply any time of day, but evening applications are preferred
- Do not use in or around Hanalei, Kaua'i (USFW Refuge)

EQUIS 720

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Beans, Dry	Anthracnose Ascochyta Blight Cercospora Leaf Blotch Downy Mildew Rust	1.37-2 pts/A	14 days	7-10 days
	<p>Specific Directions: *Seasonal limits (pints/A/year): 8 *Begin applications at first onset of disease which may occur as Early as 2 to 4 weeks before flowering *Repeat applications at 7-10 day intervals, but do not apply at less than 7 day intervals *For use only on beans to be harvested dry with pods removed. Do not apply within 14 days to harvest</p>			
Beans, Snap	Rust	1.37-3 pts/A	7 days	7 days
	<p>Specific Directions: *Seasonal limits (pints/A/year): 12 *Begin applications during Early bloom stage or when disease first threatens and repeat as necessary *Do not apply at less than 7 day intervals *Do not apply within 7 days of harvest *For resistance management of rust, alternate with another fungicide registered for bean rust control</p>			
	Botrytis Blight (Gray Mold)	3 pts/A	7 days	7 days
<p>Specific Directions: *Seasonal limits (pints/A/year): 12 *Begin applications during Early bloom stage or when disease first threatens and repeat as necessary *Do not apply at less than 7 day intervals *Do not apply within 7 days of harvest *For resistance management of rust, alternate with another fungicide registered for bean rust control</p>				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Brassica Leafy Vegetables: Broccoli, Cabbage, Cauliflower, Chinese Cabbage, Mustard Greens	Alternaria Leaf Spot Downy Mildew	1.5 pts/A	7 days	7-10 days
Specific Directions: *Use in sufficient water to obtain adequate coverage, applying by ground, air, or chemigation *Begin applications after transplants are set in field or shortly after emergence of field-seeded crop or when conditions favor disease development				
Carrots	Alternaria Leaf Blight Cercospora Leaf Spot	1.5-2.0 pts/A	0 days	0 days
Specific Directions: *Use in sufficient water to obtain adequate coverage, applying by ground, air, or chemigation *Start applications when disease threatens and repeat at 7- to 10-day intervals, but do not apply at less than 7-day intervals *Equis 720 SST may be applied the day of harvest				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Cucurbits: Cantaloupe Cucumber Gourds Honeydew Bitter melon Muskelon Pumpkin Squash Watermelon	Anthracnose Downy Mildew Target Spot	1.5- 2 pts/A	0 days	0 days
<p>Specific Directions:</p> <p>*Use in sufficient water to obtain adequate coverage, applying by ground, air, or chemigation</p> <p>*Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development</p> <p>*Repeat applications at 7-day intervals, but do not apply at less than 7-day intervals</p> <p>*Equis 720 SST may be applied the day of harvest</p> <p>Watermelon Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. Do not apply Equis 720 SST to watermelons when any of the following conditions are present: x Intense heat and sunlight x Drought conditions x Poor vine canopy x Other crop and environmental conditions which may be conducive to increased natural sunburn Do not combine Equis 720 SST with anything excepts water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use</p>				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Alternaria Leaf Blight Alternaria Leaf Spot Cercospora Leaf Spot Gummy Stem Blight/Vine Decline Powdery Mildew (Sphaerotheca only) Scab	2.0-3.0 pts/A	7 days	7 days
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water to obtain adequate coverage, applying by ground, air, or chemigation *Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development *Repeat applications at 7-day intervals, but do not apply at less than 7-day intervals *Equis 720 SST may be applied the day of harvest <p>Watermelon Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. Do not apply Equis 720 SST to watermelons when any of the following conditions are present: x Intense heat and sunlight x Drought conditions x Poor vine canopy x Other crop and environmental conditions which may be conducive to increased natural sunburn Do not combine Equis 720 SST with anything excepts water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use.</p>				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Corn, Sweet	Helminthosporium Leaf Blight Rust	0.75-2.0 pts/A	14 days	7 days
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water to obtain adequate coverage, applying by ground, air, or chemigation *Begin applications when conditions favor disease development and repeat at 7-day intervals, but do not apply at less than 7-day intervals *Under severe disease conditions, use 1.5 to 2 pints per acre *Do not apply within 14 days of harvest *Do not apply to sweet corn to be processed *Do not ensile treated corn or use as livestock forage *Do not allow livestock to graze in treated fields 			
Fruiting Vegetables: (Except Tomato) Eggplant Okra Pepino Pepper (Bell, Chili, Cooking, Pimento, Sweet) Tomatillo	Anthracnose Botrytis Blight (Gray Mold) Cercospora Leaf Spot Powdery Mildew	1.5 pts/A	3 days	7-10 days
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water to obtain adequate coverage, applying by ground, air, or chemigation *Begin applications as a foliage, flower, and fruit spray when disease is expected *Repeat applications at 7- to 10-day intervals, but do not apply at less than 7-day intervals *Do not apply within 3 days of harvest 			
Mango	Anthracnose	2.0-3.5 pts/A	21 days	7-14 days
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use 20 to 300 gallons per acre, applying by ground or air *Begin applications at Early bloom and repeat on a 7- to 14-day interval until Early fruit development, but do not apply at less than 7-day intervals *Begin the season with the 2 pint rate on a 14-day treatment interval *If disease pressure is severe, use the higher rate and shorter interval *Do not apply within 21 days of harvest 			

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Onions (Dry bulb) Garlic	Botrytis Blight (Gray Mold) Purple Blotch Botrytis Neck Rot (Suppression) Downy Mildew (Suppression)	1.0-3.0 pts/A	7 days	See label
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water to obtain thorough coverage of tops, applying by ground, air, or chemigation *Repeat applications as necessary (see table on label), but do not apply at less than 7-day intervals *Use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard (Read the label for directions) *For Suppression of neck Rot (Botrytis spp.) during storage, use a minimum of 3 weekly applications prior to lifting using 1 3/8 to 3 pints of Equis 720 SST per acre *Do not apply within 7 days of harvest 				
Papaya	Alternaria Fruit Spot Anthracnose Stem End Rot	1.5-3.0 pts/A	0 days	14 days
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use sufficient water to obtain adequate coverage of fruit and leaves, applying by ground only *Begin treatment when conditions favor development of disease, and continue treatments at 14-day intervals until weather conditions no longer favor disease development *Do not apply at less than a 14-day interval *Equis 720 SST may be applied the day of harvest 				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Peanut	Early Leaf Spot (Cercospora) Late Leaf Spot (Cercosporidium) Pepper Spot	1.0-1.5 pts/A	14 days	14 days
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting, using ground, air, or chemigation *Repeat at 14-day intervals, but do not apply at less than 14-day intervals. *When conditions favor Late Leaf Spot or when rust or web blotch occur, apply 1.5 pints per acre at 14-day intervals for the remainder of the season *Alternate chemigation applications with ground or aerial applications *Use 1.5 pints if applying by chemigation *Do not apply within 14 days of harvest *Do not allow livestock to graze in treated areas *Do not feed hay or threshings from treated fields to livestock 			
	Rust Web Blotch	1.5 pts/A	14 days	14 days
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting, using ground, air, or chemigation *Repeat at 14-day intervals, but do not apply at less than 14-day intervals. *When conditions favor Late Leaf Spot or when rust or web blotch occur, apply 1.5 pints per acre at 14-day intervals for the remainder of the season *Alternate chemigation applications with ground or aerial applications *Use 1.5 pints if applying by chemigation *Do not apply within 14 days of harvest *Do not allow livestock to graze in treated areas *Do not feed hay or threshings from treated fields to livestock 				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Tomato	Foliage: Early Blight Botrytis Blight (Gray Mold) Gray Leaf Spot Late Blight Septoria Leaf Spot Target Spot	1.37-2.0 pts/A	0 days	7-10 days
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water to obtain adequate coverage, applying by ground, air, or chemigation *Begin applications when dew or rain occurs and disease threatens *Apply every 7 to 10 days for foliage diseases *For fruit diseases, begin at fruit set and apply every 7 to 14 days *Do not apply at less than 7-day intervals *Use the highest rate and shortest interval when disease is severe *Equis 720 SST may be applied the day of harvest 				
	Alternaria Fruit Rot (Black Mold) Anthracnose Botrytis Blight (Gray Mold) Late Blight Fruit Rot Rhizoctonia Fruit Rot	2.0-2.75 pts/A	0 days	7-14 days
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Use in sufficient water to obtain adequate coverage, applying by ground, air, or chemigation *Begin applications when dew or rain occurs and disease threatens *Apply every 7 to 10 days for foliage diseases *For fruit diseases, begin at fruit set and apply every 7 to 14 days *Do not apply at less than 7-day intervals *Use the highest rate and shortest interval when disease is severe *Equis 720 SST may be applied the day of harvest 				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Yam	Anthracnose	1.0-1.25 pts/A	7 days	10-14 days
<p>Specific Directions: *Use in sufficient water to obtain adequate coverage, applying by ground, air, or chemigation *Begin applications when disease first threatens *Repeat applications As needed at 10- to 14- day intervals, but do not apply at less than 10-day intervals. *Do not apply within 7 days of harvest</p>				

KOCIDE 3000 **

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Avocado	Anthracnose Blotch Scab	3.5-5.25 lb/A	0 days	14-30 days
	Specific Directions: *Apply when bloom buds begin to swell and continue application at 14-30 day intervals for five to six applications *Use higher rates when conditions favor disease *Maximum annual rate/acre: 63 lb.			
Banana	Sigatoka (Black and Yellow)	0.75 lb/A	0 days	7-14 days
	Specific Directions: *Maximum annual rate/acre: 63 lb.			
Beans (Dry, Green)	Brown Spot Common Blight	0.5-1.25 pts/A	0 days	7-14 days
	Specific Directions: *Read the label. *Use the higher rate for more severe disease *For protective sprays, make first application when plants are 6 inches high; repeat on a 7-14 day schedule depending upon local conditions			
Cacao	Black Pod	0.75-3.75 lb/A	0 days	14-21 days
	Specific Directions: *Begin applications at the start of the rainy season and continue while infection conditions persist *Apply .75 to 2 pounds at 14 to 21 day intervals if needed depending on disease severity *For drier areas, make two to four applications using 2.5 to 3.75 pounds per acre according to disease incidence and planting density *Maximum annual rate/acre: 52.5 lb.			

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Carrot	Alternaria Leaf Spot Cercospora Leaf Spot	0.75 - 1.5 lb/A	0 days	7-14 days
	<p>Specific Directions: *Begin applications when disease first threatens and repeat at 7- to 14- day intervals if needed depending on disease severity *Maximum annual rate/acre: 16.7 lb.</p>			
Citrus: Grapefruit Lemon Lime Orange Tangerine	Melanose Scab	1.75 - 5 lb/A	0 days	7 days
	<p>Specific Directions: *Apply as pre-bloom and post-bloom sprays *Use the higher rates when conditions favor disease *Minimum retreatment interval is 7 days *Maximum annual rate/acre: 42 lb.</p>			
	Alternaria Brown Spot	1.75 - 3.5 lb/A	0 days	7 to 21 days
<p>Specific Directions: *On susceptible varieties apply when the first spring flush appears and each flush thereafter *Application to fruit should start after two thirds of the petals have fallen and be repeated on a 7 to 21 day schedule if needed *Use the higher rates when conditions favor disease *Maximum annual rate/acre: 42 lb.</p>				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Phytophthora Brown Rot Septoria Spot	1.75-3.5 lb/A	0 days	7 days
<p>Specific Directions: *Begin application in fall before or just after the first rain and continue if needed *For Brown Rot only, apply to skirts of trees to a height of at least 4 feet *For control of Septoria Spot or where fruit have already been infected with Brown Rot, apply to entire tree *Apply also to bare ground one Foot beyond skirt *Use the higher rates when conditions favor disease *Minimum retreatment interval is 7 days *Maximum annual rate/acre: 42 lb.</p>				
	Phytophthora Foot Rot	0.5 lb/A	0 days	1 year
<p>Specific Directions: *Mix with 1 quart of water, "Tre-Hold" or Latex paint. Paint trunks of trees from the soil surface to the lowest scaffold limbs *Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection *Treatment serves as protection for up to 1 year, but does not cure existing infections *NOTE: Areas where microjet or low volume irrigation hit the tree trunk may require retreatment due to wash off. *Maximum annual rate/acre: 42 lb.</p>				
	Citrus Canker (Suppression)	1 - 2.5 lb/A	0 days	7 days
<p>Specific Directions: *Spray flushes 7 to 14 days after shoots begin to grow *Young fruit may require an additional application *Number and timing of applications will be dependent upon disease pressure *Under heavy pressure, spray each flush of new growth *Minimum retreatment interval is 7 days *Maximum annual rate/acre: 42 lb.</p>				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Black Spot	1 - 3 lb/A	0 days	7-21 days
<p>Specific Directions: *Begin treatment prior to or when disease first appears and repeat every 7 to 21 days if needed *Use the higher rates and shorter spray intervals when conditions favor disease *Minimum retreatment interval is 7 days *Maximum annual rate/acre: 42 lb.</p>				
Corn (Field, Sweet)	Bacterial Stalk Rot	0.5 - 1.75 lb/A	0 days	7-10 days
<p>Specific Directions: *Begin treatment when disease first appears and repeat every 7 to 10 days if needed *Use the higher rates and shorter spray intervals when conditions favor disease *Maximum annual rate/acre: 14 lb.</p>				
<p>Crucifers: Broccoli Cabbage Chinese Cabbage Cauliflower Collard Greens Mustard Greens Turnip Kale</p>	<p>Black Leaf Spot (Alternaria) Black Rot (Xanthomonas) Downy Mildew</p>	0.5 - 0.75 lb/A	0 days	7-10 days
<p>Specific Directions: *Begin application after transplants are set in the field, or shortly after emergence of field seeded crops or when conditions favor disease development *Apply at 7- to 10-day intervals if needed *Use the higher rates when conditions favor disease *NOTE: Reddening of older leaves may occur on broccoli and a flecking of wrapper leaves may occur on cabbage *Maximum annual rate/acre: 8.8 lb.</p>				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Cucurbits: Cantaloupe Cucumber Honeydew Muskmelon Pumpkin Squash Watermelon	Alternaria Leaf Spot Angular Leaf Spot Anthracnose Downy Mildew Gummy Stem Blight Powdery Mildew Watermelon Bacterial Fruit Blotch (Suppression)	0.5 - 1.25 lb/A	0 days	5-7 days
<p>Specific Directions: *Begin applications prior to disease development and continue while conditions are favorable for disease development *Repeat at 5- to 7-day intervals if needed *Use the higher rates when conditions favor disease *NOTE: Crop injury may occur from application at higher rates and shorter intervals. Discontinue use if injury occurs *Maximum annual rate/acre: 17.5 lb.</p>				
Eggplant	Alternaria Blight Anthracnose	0.75 - 1.5 lb/A	0 days	7-10 days
<p>Specific Directions: *Begin applications prior to development of disease symptoms *Repeat sprays at 7- to 10-day intervals if needed depending on disease severity *Maximum annual rate/acre: 26.3 lb.</p>				
Guava	Anthracnose Red Algae	1.25 - 2.0 lb/A	0 days	7 days
<p>Specific Directions: *Make initial application just before flowering and repeat on a weekly schedule until just before harvest *Apply in sufficient water for thorough coverage *Use the higher rates for severe disease *Maximum annual rate/acre: 16.4 lb.</p>				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Lettuce	Downy Mildew	0.75 - 1.5 lb/A	0 days	5-10 days
	<p>Specific Directions: *Begin applications when disease symptoms first appear or when conditions favor disease development *Repeat at 5- to 10-day intervals if needed depending on disease severity *NOTE: Determine if there is varietal sensitivity prior to use *Injury may occur to sensitive lettuce varieties and under adverse weather conditions *Discontinue use if injury occurs *Maximum annual rate/acre: 26.6 lb.</p>			
Mango	Anthracnose	2 - 6 lb/A	0 days	7 days
	<p>Specific Directions: *Apply at 7 day intervals after fruit set until harvest *Use the higher rates when rainfall is heavy and disease pressure is high *Maximum annual rate/acre: 160 lb.</p>			
Okra	Anthracnose Bacterial Leaf Spot Leaf Spot Pod Spot Powdery Mildew	0.75 - 1.75 lb/A	0 days	5-10 days
	<p>Specific Directions: *Begin treatment when disease first threatens and repeat every 5- to 10- days if needed depending on disease severity *Use the higher rates and shorter spray intervals when conditions favor disease *Maximum annual rate/acre: 17.5 lb.</p>			
Onion Garlic	Bacterial Blight Downy Mildew Purple Blotch	0.75 - 1.5 lb/A	0 days	7-10 days
	<p>Specific Directions: *Begin when plants are 4 to 6 inches high and repeat at 7- to 10-day intervals if needed depending on disease severity *Can cause phytotoxicity to leaves *Maximum annual rate/acre: 20 lb.</p>			

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Peanut	Cercospora Leaf Spot	0.75 - 1.25 lb/A	0 days	7-14 days
	Specific Directions: *Begin spraying at 35 to 40 days after planting or when disease symptoms first appear and repeat at 7- to 14-day intervals if needed *Reduce sprays to 7 day intervals during humid weather *Use the higher rates when conditions favor disease *Flowable sulfur may be added *Maximum annual rate/acre: 15.8 lb.			
Pepper	Anthracnose Bacterial Spot Cercospora Leaf Spot	0.75 - 1.25 lb/A	0 days	3-10 days
	Specific Directions: *Begin applications when conditions first favor disease development and repeat at 3- to 10-day intervals if needed depending on disease severity *Use the higher rates when conditions favor disease *Maximum annual rate/acre: 39.5 lb.			
Papaya	Anthracnose	1.75 - 4.25 lb/A	0 days	7 days
	Specific Directions: *Apply before disease appears. Apply at 7-day intervals if needed *The addition of an approved spreader is desirable *Use the higher rates when conditions favor disease *Maximum annual rate/acre: 70.7 lb.			
Parsley	Bacterial Blight (Pseudomonas sp.)	1.25 - 2.0 lb/A	0 days	10 days
	Specific Directions: *Begin applications when plants are first established in the field and repeat at 10 day intervals if needed depending on disease severity and environmental conditions *Maximum annual rate/acre: 6.7 lb.			

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Tomato	Anthracnose Bacterial Speck Bacterial Spot Botrytis Blight (Gray Mold) Late Blight Septoria Leaf Spot	0.75 - 1.75 lb/A	0 days	3-10 days
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Begin applications when disease first threatens and repeat at 3- to 10- day intervals if needed depending on disease severity *Use the higher rates when conditions favor disease *Maximum annual rate/acre: 58 lb. (Processing), 26.7 lb. (Fresh market) 			

LANNATE LV

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Avocado	Leafrollers Loopers	1.5-3 pts/A	1 day	5-7 days
	Specific Directions: *Read the label *Do not apply more than 3 pints of LANNATE®LV/acre (A)/crop *Do not make more than 2 applications/crop			
Beans	Beetles Leafhoppers	0.75-3 pts/A	See label	5-7 days
	Specific Directions: *Read the label *Do not apply more than 15 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop			
	Cutworms	1.5 pts/A	See label	5-7 days
	Specific Directions: *Read the label *Do not apply more than 15 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop			
	Armyworms Aphids Caterpillars Loopers Thrips	1.5-3 pts/A	See label	5-7 days
Specific Directions: *Read the label *Do not apply more than 15 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Broccoli	Diamondback Moths Loopers	1.5-3 pts/A	3 days	2 days
	Specific Directions: *Read the label *Do not apply more than 21 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop *Add a wetting agent to improve coverage			
	Caterpillars	0.75-3 pts/A	3 days	2 days
	Specific Directions: *Read the label *Do not apply more than 21 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop *Add a wetting agent to improve coverage			
Cabbage	Diamondback Moths Loopers	1.5-3 pts/A	1 day	5-7 days
	Specific Directions: *Read the label *Do not apply more than 24 pints of LANNATE®LV/acre (A)/crop *Do not make more than 15 applications/crop *Add a wetting agent to improve coverage			
	Caterpillars	0.75-3 pts/A	1 day	2 days
	Specific Directions: *Read the label *Do not apply more than 24 pints of LANNATE®LV/acre (A)/crop *Do not make more than 15 applications/crop *Add a wetting agent to improve coverage			
	Armyworms Cutworms	1.5 pts/A	1 day	2 days
Specific Directions: *Read the label *Do not apply more than 24 pints of LANNATE®LV/acre (A)/crop *Do not make more than 15 applications/crop *Add a wetting agent to improve coverage				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Carrot	Leafhoppers	1.5-3 pts/A	1 day	5-7 days
	Specific Directions: *Read the label *Do not apply more than 21 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop			
	Armyworms Cutworms	0.75-1.5 pts/A	1 day	5-7 days
	Specific Directions: *Read the label *Do not apply more than 21 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop			
Cauliflower	Caterpillars	0.75-3 pts/A	3 days	2 days
	Specific Directions: *Read the label. *Do not apply more than 24 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop *Add a wetting agent to improve coverage			
	Diamondback Moths Loopers	1.5-3 pts/A	3 days	2 days
	Specific Directions: *Read the label *Do not apply more than 24 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop *Add a wetting agent to improve coverage			
	Cutworms	1.5 pts/A	3 days	2 days
	Specific Directions: *Read the label *Do not apply more than 24 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop *Add a wetting agent to improve coverage			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Chinese Cabbage (Napa, Bok Choy)	Armyworms Loopers	1.5-3 pts/A	10 days	5-7 days
	Specific Directions: *Read the label *Do not apply more than 24 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop *Minimum of 25 gallons water per acre			
Corn, Field	Aphids Armyworms Beetles Caterpillars Flea Beetles	0.75-1.5 pts/A	See label	5-7 days
	Specific Directions: *Read the label *Do not apply more than 7.5 pints of LANNATE®LV/acre (A)/crop *Do not make more than 5 applications/crop			
	Cutworms	1.5 pts/A	See label	5-7 days
	Specific Directions: *Read the label *Do not apply more than 7.5 pints of LANNATE®LV/acre (A)/crop *Do not make more than 5 applications/crop			
Corn, Sweet	Aphids Armyworms Beetles Caterpillars Flea Beetles	0.75-1.5 pts/A	See label	1 day
	Specific Directions: *Read the label *Do not apply more than 21 pints of LANNATE®LV/acre (A)/crop *Do not make more than 28 applications/crop *Certain hybrid varieties of corn are susceptible to methomyl injury *Treat a small area to determine crop safety before full scale spraying			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Cutworms	1.5 pts/A	See label	1 day
Specific Directions: *Read the label *Do not apply more than 21 pints of LANNATE®LV/acre (A)/crop *Do not make more than 28 applications/crop *Certain hybrid varieties of corn are susceptible to methomyl injury *Treat a small area to determine crop safety before full scale spraying				
Cucumber	Aphids Armyworms Caterpillars Cucumber Beetles Cutworms Flea Beetles Melonworms	1.5-3 pts/A	See label	5-7 days
Specific Directions: *Read the label *Do not apply more than 18 pints of LANNATE®LV/acre (A)/crop *Do not make more than 12 applications/crop				
Eggplant	Aphids	0.75-3 pts/A	5 days	5-7 day
Specific Directions: *Read the label *Do not apply more than 15 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop				
	Armyworms Caterpillars	1.5-3 pts/A	5 days	5-7 days
Specific Directions: *Read the label *Do not apply more than 15 pints of LANNATE®LV/acre (A)/crop *Do not make more than 10 applications/crop				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Garlic	Armyworms	1.5-3 pts/A	7 days	5-7 days
	Specific Directions: *Read the label *Do not apply more than 9 pints of LANNATE®LV/acre (A)/crop *Do not make more than 6 applications/crop *Add a wetting agent to improve coverage			
Horseradish	Aphids Thrips	1.5 pts/A	65 days	5-7 days
	Specific Directions: *Read the label *Do not apply more than 6 pints of LANNATE®LV/acre (A)/crop *Do not make more than 4 applications/crop			
Leafy Green Vegetables: Mustard Greens Parsley	Armyworms Caterpillars Diamondback Moths Loopers	1.5-3 pts/A	10 days	5-7 days
	Specific Directions: *Read the label *Do not apply more than 12 pints of LANNATE®LV/acre (A)/crop *Do not make more than 8 applications/crop			
Lettuce (Head, Leaf)	Aphids Armyworms Loopers Caterpillars Leafhoppers Thrips	1.5-3 pts/A	See label	2 days
	Specific Directions: *Read the label Lettuce (Head varieties): *Do not apply more than 21 pints of LANNATE®LV/acre (A)/crop *Do not make more than 12 applications/crop Lettuce (Leaf varieties): *Do not apply more than 12 pints of LANNATE®LV/acre (A)/crop *Do not make more than 6 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Cutworms	1.5 pts/A	See label	2 days
<p>Specific Directions: *Read the label</p> <p>Lettuce (Head varieties): *Do not apply more than 21 pints of LANNATE®LV/acre (A)/crop *Do not make more than 12 applications/crop</p> <p>Lettuce (Leaf varieties): *Do not apply more than 12 pints of LANNATE®LV/acre (A)/crop *Do not make more than 6 applications/crop</p>				
<p>Melons:</p> <p>Cantaloupe Honeydew Muskmelon Pepino Watermelon</p>	<p>Aphids Armyworms Loopers Cucumber Beetles Cutworms Caterpillars Flea Beetles Melonworms</p>	1.5-3 pts/A	See label	5-7 days
<p>*Read the label * Do not apply more than 18 pints of LANNATE®LV/acre (A)/crop *Do not make more than 12 applications/crop</p>				
<p>Onion, Green</p>	<p>Armyworms Cutworms Thrips</p>	1.5-3 pts/A	7 days	5-7 days
<p>Specific Directions: *Read the label *Do not apply more than 21 pints of LANNATE®LV/acre (A)/crop *Do not make more than 8 applications/crop *Add wetting agent to improve coverage</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Peanut	Armyworms Caterpillars Cutworms Leafhoppers Loopers Thrips	0.75-3 pts/A	21 days	5-7 days
	Specific Directions: *Read the label *Do not apply more than 12 pints of LANNATE®LV/acre (A)/crop *Do not make more than 8 applications/crop			
Pepper (Bell, Hot)	Aphids Armyworms Loopers Caterpillars Cutworms	0.75-3 pts/A	3 days	5-7 days
	Specific Directions: *Read the label *Do not apply more than 12 pints of LANNATE®LV/acre (A)/crop *Do not make more than 8 applications/crop			
Squash, Summer Bitter melon Gourds Kondot	Aphids Armyworms Loopers Cucumber Beetles Cutworms Caterpillars Flea Beetles Melonworms	1.5-3 pts/A	See label	5-7 days
	Specific Directions: *Read the label *Do not apply more than 18 pints of LANNATE®LV/acre (A)/crop *Do not make more than 12 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Tomato	Aphids Armyworms Caterpillars Cutworms Loopers	0.75-3 pts/A	1 day	5-7 days
	Specific Directions: *Read the label *Do not apply more than 7.2 lbs. a.i./ A /crop *Do not make more than 16 applications/crop			

MALATHION 5 EC

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Avocado	Loopers Scales Thrips	7.5 pts/a	7 days	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Broccoli	Aphids Caterpillars Loopers	1-2 pts/A	2 days	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Cabbage	Aphids Caterpillars Loopers Diamondback Moths	1-2 pts/A	7 days	As needed
	Specific Directions: *Read the label *Do not make more than 6 applications/crop			
Carrot	Aphids	1.5-2 pts/A	7 days	As needed
	Specific Directions: *Read the label			
	Leafhoppers	1.5-2 pts/A	7 days	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Cauliflower	Aphids Diamondback Moths	1-2 pts/A	7 days	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Citrus: Grapefruit Lemon Lime Orange Tangerine	Mites Scales Thrips	2.4 pts/A	7 days	As needed
	Specific Directions: *Read the label *Do not apply when trees are in bloom *Do not make more than 3 applications/crop			
Corn, Sweet	Beetles	1.6 pts/A	5 days	As needed
	Specific Directions: *Read the label *Injury may occur in the whorl or to the silks *Do not make more than 2 applications/crop			
Cucumber	Aphids Caterpillars Beetles Cucumber Beetles Leafminers Spider Mites	1.5-2.8 pts/A	1 day	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Eggplant	Aphids Spider Mites	1.5-2.5 pts/A	3 days	As needed
	Specific Directions: *Read the label *Do not make more than 4 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Garlic	Aphids Thrips	1.5-2.5 pts/A	3 days	As needed
	Specific Directions: *Read the label *Do not make more than 3 applications/crop			
Lettuce, Leaf	Aphids Leafhoppers Spider Mites	2-3 pts/A	14 days	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
	Loopers Scales Thrips	2.5-3 pts/A	14 days	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Melons: Cantaloupe Honeydew Muskmelon	Aphids Cucumber Beetles Leafminers Leafhoppers Spider Mites	1.5-2 pts/a	1 day	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Okra	Aphids	1.5 pts/A	1 day	As needed
	Specific Directions: *Read the label * Do not make more than 5 applications/crop			
	Beetles	1.9 pts/A	1 day	As needed
	Specific Directions: *Read the label *Do not make more than 5 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Papaya	Aphids Mealybugs	1.5-2 pts/A	1 day	As needed
	Specific Directions: *Read the label *Do not make more than 8 applications/crop			
Parsley	Aphids	1.5-2 pts/A	21 days	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Pepper (Bell, Hot)	Aphids Caterpillars	1-2.5 pts/A	3 days	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Pineapple	Mealybugs	1 pts/100 gals water or 8 pts/A	7 days	As needed
	Specific Directions: *Read the label *Do not make more than 3 applications/crop			
Pumkin	Aphids Caterpillars Cucumber Beetles Leafhoppers Spider Mites	1.5 pts/A	1 day	As needed
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Radish Horseradish	Aphids	1.5 pts/A	7 days	As needed
	Specific Directions: *Read the label *Do not make more than 3 applications/crop			
Squash, Summer	Aphids Cucumber Beetles Caterpillars Leafminers Spider Mites	2-2.8 pts/A	1 day	As needed
	Specific Directions: *Read the label. *Do not make more than 3 applications/crop			
Squash, Winter	Aphids Cucumber Beetles Caterpillars Leafminers Spider Mites	1.6 pts/A	1 day	As needed
	Specific Directions: *Read the label. *Do not make more than 3 applications/crop			
Sweet Potato	Leafhoppers Leafminers	1.5-2.5 pts/A	0 days	As needed
	Specific Directions: *Read the label. *Do not make more than 2 applications/crop			
Tomato	Aphids Spider Mites	1.5 pts/A	1 day	As needed
	Specific Directions: *Read the label			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Armyworms Caterpillars	2.5 pts/A	1 day	As needed
<p>Specific Directions: *Read the label *Do not make more than 4 applications/crop</p>				
Watermelon	Aphids Cucumber Beetles Leafminers Leafhoppers Spider Mites	1.5-2.5 pts/A	1 day	As needed
<p>Specific Directions: *Read the label *Do not make more than 2 applications/crop</p>				

MALATHION 8 AQUAMUL

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Avocado	Loopers Scales Thrips	4-4.7pts/A	7 days	30 days
	Specific Directions: *Read the label *Do not make more than 2 applications/acre per calendar year			
Broccoli	Aphids Caterpillars Loopers	1.25pts/A	2 days	7 days
	Specific Directions: *Read the label *Do not make more than 6 applications/acre per calendar year			
Cabbage	Aphids Caterpillars Diamondback Moths Loopers	1.25pts/A	7 days	See label
	Specific Directions: *Read the label *Do not make more than 6 applications/acre per calendar year			
Carrot	Aphids Leafhoppers	1-1.25 pts/A	7 days	7 days
	Specific Directions: *Read the label *Do not make more than 2 applications/acre per calendar year			
Citrus: Grapefruit Lemon Lime Orange Tangerine	Scales	1.5pts/A	7 days	30 days
	Specific Directions: *Read label *Do not apply when trees are in bloom *Do not make more than 3 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Corn, Sweet	Beetes	1 pt/A	5 days	5 days
	Specific Directions: *Read the label *Injury may occur in the whorl or to the silks *Do not make more than 2 applications/crop			
Cucumbers	Aphids Beetles Caterpillars Cucumber Beetles Leafhoppers Spider Mites	1-1.75 pts/A	1 day	7 days
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Eggplant	Aphids Spider Mites	0.75-1.56 pts/A	3 day	5 days
	Specific Directions: *Read the label *Do not make more than 4 applications/crop			
Garlic	Aphids Thrips	1-1.5 pts/A	3 days	7 days
	Specific Directions: *Read the label *Do not make more than 3 applications/crop			
Lettuce, Leaf	Aphids Leafhoppers	1.25-1.88 pts/A	14 days	6 days
	Specific Directions: *Read label *Do not make more than 3 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Loopers Spider Mites	1.88 pts/A	14 days	6 days
Specific Directions: *Read label *Do not make more than 3 applications/crop				
Melons: (Other than Watermelon)	Aphids Cucumber Beetles Leafhoppers Leafminers Spider Mites	1 pts/A	1 day	7 days
Specific Directions: *Read label *Do not make more than 2 applications/crop				
Okra	Aphids Beetles	1.2 pts/A	1 day	7 days
Specific Directions: *Read the label *Do not make more than 5 applications/crop				
Papaya	Aphids Mealybugs	1.25 pts/A	1 day	3 days
Specific Directions: *Read the label *Do not make more than 8 applications/crop				
Parsley	Aphids	1-1.5 pts/A	7 days	7 days
Specific Directions: *Read label *Do not make more than 2 applications/crop				
Peppers	Aphids	0.75-1.5 pts/A	3 days	5 days
Specific Directions: *Read the label *Do not make more than 2 applications/crop				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Pineapple	Mealybugs	2 pts/A	7 days	7 days
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Pumpkin	Aphids Caterpillars Cucumber Beetles Leafhoppers Spider Mites Thrips	1 pt/A	1 day	7 days
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Radish Horseradish	Aphids	1 pt/A	7 days	7 days
	Specific Directions: *Read the label *Do not make more than 3 applications/crop			
Squash, Summer	Aphids Caterpillars Cucumber Beetles Spider Mites Thrips	1.75 pts/A	1 day	7 days
	Specific Directions: *Read the label *Do not make more than 3 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Squash, Winter	Aphids Caterpillars Cucumber Beetles Spider Mites Thrips	1 pt/A	1 day	7 days
	Specific Directions: *Read the label *Do not make more than 3 applications/crop			
Sweet Potato	Leafhoppers	1-1.56 pts/A	0 days	7 days
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
	Leafminers	1 3/4 pts/A	0 days	7 days
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			
Tomato	Aphids Spider Mites	1-1.56 pts/A	1 day	5 days
	Specific Directions: *Read the label *Do not make more than 4 applications/crop			
	Armyworms Caterpillars	1.56 pts/A	1 day	5 days
	Specific Directions: *Read the label *Do not make more than 2 applications/crop			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Watermelon	Aphids Cucumber Beetles Leafhoppers Leafminers Spider Mites	1.5 pts/A	1 day	7 days
	Specific Directions: *Read label *Do not make more than 2 applications/crop			

MANZATE PRO STICK

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Banana	Sigatoka	2-3 lb/A	0 days	14-21 days
	<p>Specific Directions: *Apply when leaves first appear and repeat every 14 to 21 days or as required *Use sufficient water to provide adequate coverage *Do not apply more than 30 lbs (22.5 lbs active) per acre per growing cycle *Minimum preharvest interval 0 days</p>			
Broccoli	Alternaria Leaf Spot Downy Mildew	1.6-2.1 lb/A	7 days	7-10 days
	<p>Specific Directions: *Begin applications prior to disease development and when conditions are favorable for disease development *Apply at 7 to 10-day intervals, if needed *Use higher rates when conditions favor disease *Do not apply more than 12.8 lbs product (9.6 lbs)/A/season *Minimum retreatment interval is 7 days *Minimum preharvest interval is 7 days *Do not apply this product with a U-boom device</p>			
Cabbage	Alternaria Leaf Spot Downy Mildew	1.6-2.1 lb/A	7 days	7-10 days
	<p>Specific Directions: *Begin applications prior to disease development and when conditions are favorable for disease development *Apply at 7 to 10-day intervals, if needed *Use higher rates when conditions favor disease *Do not apply more than 12.8 IDS product (9.6 lbs)/A/season *Minimum retreatment interval is 7 days *Minimum preharvest interval is 7 days *Do not apply this product with a U-boom device</p>			

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Corn, Sweet	Common Rust Helminthosporium Leaf Blight Gray Leaf Spot	1.5 lb/A	7 days	4-7 days
	<p>Specific Directions: *Use sufficient water for thorough coverage *Start applications when disease first appears and repeat at 4 to 7 day intervals *Do not apply within 7 days of harvest *Do not apply more than 22.5 lbs (17 lbs active) per acre per crop east of the Mississippi and AR and LA *Do not apply more than 7.5 lbs (5.6 lbs active) per acre per crop west of the Mississippi excepts AR and LA *Do not apply to sweet corn in home gardens</p>			
Corn, Field	Common Rust Helminthosporium Leaf Blight Gray Leaf Spot	1.5 lb/A	40 days	4-7 days
	<p>Specific Directions: *Do not apply within 40 days of harvest *Do not apply more than 15 lbs (11.25 lbs active) per acre per crop</p>			
Cucumber	Anthracnose Cercospora Leaf Spot Downy Mildew Gummy Stem Blight Scab	2-3 lb/A	5 days	7-10 days
	<p>Specific Directions: *Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals *Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces *Do not apply within 5 days of harvest *Do not apply more than 24 lbs (18 lbs active) per acre per crop</p>			

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
<p>Cucurbits: Chinese Wax Gourd Gourd Muskmelon Pumpkin Squash, Summer Squash, Winter Watermelon</p>	<p>Alternaria Leaf Spot Anthracnose Cercospora Leaf Spot Downy Mildew Gummy Stem Blight Scab</p>	<p>2-3 lb/A</p>	<p>5 days</p>	<p>7-10 days</p>
<p>Specific Directions: *Start applications when the plants are in the two-leaf stage and repeat at 7- to 10- day intervals *Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces *For aerial applications, the minimum spray volume is 2 gallons per acre *Do not apply more than 25.6 lbs. product (19.2 lbs. active ingredient) per acre per year *Do not apply more than 8 applications per year *Do not apply within 5 days of harvest *Note: Some cantaloupe varieties (i.e. Harvest Queen, Gold Star, Super Star, Sweet and Early, and Saticoy) are sensitive to Manzate Pro-Stick fungicide *Consult UOG Cooperative Extension Service Specialist prior to use</p>				
<p>Lettuce (Head, Leaf)</p>	<p>Anthracnose Downy Mildew</p>	<p>1.6-2.1 lb/A</p>	<p>See label</p>	<p>7-10 days</p>
<p>Specific Directions: *Begin applications prior to disease development and when conditions are favorable for disease development *Apply at 7 to 10-day intervals, if needed *Use higher rates when conditions favor disease *Remove residues from head lettuce by stripping and trimming *For all states: minimum retreatment interval is 7 days *Do not apply this product with a U-boom device</p>				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
<p>Melon: Cantaloupe Honeydew Watermelon</p>	<p>Alternaria Leaf Spot Anthrachnose Downy Mildew Gummy Stem Blight Cercospora Leaf Spot</p>	<p>2-3 lb/A</p>	<p>5 days</p>	<p>7-10 days</p>
<p>Specific Directions: *Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals *Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces *Some varieties are sensitive to MANZATE PRO-STICK fungicide *Consult UOG Cooperative Extension Service Specialist prior to use *Do not apply within 5 days of harvest *Do not apply more than 24 lbs (18 lbs active) per acre per season</p>				
<p>Onion (Dry bulb) Garlic Shallot</p>	<p>Botrytis Blight (Gray Mold) Downy Mildew Neck Rot Purple Blotch</p>	<p>3 lb/A</p>	<p>7 days</p>	<p>7 days</p>
<p>Specific Directions: *Follow a protective spray schedule starting when diseases are first reported in the area and repeat at 7 day intervals throughout the season *Do not apply within 7 days of harvest *Do not apply to exposed bulbs *Do not apply more than 30 lbs (22.5 lbs active) per acre per crop</p>				

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
<p>Papaya</p>	<p>Anthracnose (Colletotricum) Phytophthora Fruit Rot Black Spot (Cercospora)</p>	<p>2.0-2.5 lb/A (minimum 50 gals per acre)</p>	<p>0 days</p>	<p>14 to 21 days</p>
	<p>Specific Directions: *Begin at flowering; treat central column crown, blossom area and developing fruit. Repeat at 14 to 21 day intervals *Do not use more than 37 lbs (28 lbs active) per acre per year, i.e. do not make more than 14 applications at the maximum use rate per year *Minimum pre-harvest interval 0 days</p>			
<p>Peanut</p>	<p>Ascochyta Web Blotch Cercospora Leaf Spot Rust</p>	<p>1-2 lb/A</p>	<p>14 days</p>	<p>7-14 days</p>
	<p>Specific Directions: *Start application when disease first appears or is reported in area *Repeat sprays at 7 to 14 day intervals *Reduce sprays to a 7 day interval during humid weather *Do not apply within 14 days of harvest *Do not use more than 16 lbs (12 lbs active) per acre per crop *Do not feed treated vines to livestock</p>			
<p>Pepper</p>				
	<p>Specific Directions: *See label</p>			
<p>Squash, Summer</p>	<p>Downy Mildew</p>	<p>2-3 lb/A</p>	<p>5 days</p>	<p>7-10 days</p>
	<p>Specific Directions: *Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals *Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces *Do not apply within 5 days of harvest *Do not apply more than 24 lbs (18 lbs active) per acre per crop</p>			

CROP	DISEASE(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Tomato				
	Specific Directions: *See label			
Tropical Fruits: Mango Star apple	Anthracnose Phytophthora Fruit Rot Black Spot (Cercospora)	2-2.5 lb/A	0 days	14-21 days
Specific Directions: *Start applications at flowering and continue at 14- to 21-day intervals *Direct spray to crown and blossom area *Use 20 to 100 gallons water per acre *Do not apply more than 37.3 lbs. product (28 lbs. active ingredient) per acre per year *Do not apply more than 14 applications per year *Applications may be made up to the day of harvest				

NEEM OIL **

Fungicide/Insecticide/Miticide/Nematicide
OMRI Listed

NEEM OIL **

See label

PERM-UP 3.2 EC

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Avocado	Caterpillar Leafhoppers Loopers Scales Scale Crawlers Thrips Whiteflies	8 oz/A	7 days	7-10 days
	Specific Directions: *Do not apply more than 32 oz. product (0.8 lbs.. a.i.) per acre per season *Apply when insects first appear and repeat as needed to provide control			
Brassica Leafy Vegetables: Broccoli Cauliflower	Aphids (Suppression) Armyworms Diamondback Moths Loopers Plant Bugs Thrips	2-4 oz/A	1 day	5 days
	Specific Directions: *Maximum rate for broccoli is 8 oz. product (0.2. lbs.. a.i.)/A *Do not apply more than 0.8 lbs.. a.i. per acre per season for broccoli *Do not apply more than 0.4 lbs.. a.i. per acre per season for cauliflower *Apply when insects first appear and repeat as needed to provide control			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Cabbage Chinese Cabbage	Armyworms Cutworms Flea Beetles	4-8 oz/A	1 day	5-10 days
	Specific Directions: *Apply every 5-10 days or as needed *Do not apply more than 16 oz. Product (0.4 lbs.. a.i.) per acre per season (32 oz. product/0.8 lbs.. a.i. in Hawaii)			
	Aphids (Suppression) Diamondback Moths Loopers	2-8 oz/A	1 day	5-10 days
Specific Directions: *Apply every 5-10 days or as needed *Do not apply at intervals less than 5 days *Do not apply more than 16 oz. Product (0.4 lbs.. a.i.) per acre per season (32 oz. product/0.8 lbs.. a.i. in Hawaii)				
Corn, Sweet	Armyworms Cutworms Flea Beetles Leafhoppers	4-8 oz/A	1 day	3-5 days
Specific Directions: *Do not apply more than 32 oz. product (0.8 lbs.. a.i.) per acre per season *Apply every 3 to 5 days or as needed *Do not apply at intervals less than 3 days *Do not apply within 1 day of harvest				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Cucurbits: Cantaloupe Cucumber Gourd Bitter melon Honeydew Muskmelon Pumpkin Squash (Summer, Winter) Watermelon	Aphids Leafminers Squash Bugs	8 oz/A	0 days	7 days
	Specific Directions: *Apply by air or ground equipment. By air, apply in a minimum of 4 gallons of water per acre *By ground, apply in a minimum of 20 gallons of water per acre *Use sufficient water to obtain full coverage of foliage			
	Cucumber Beetles Cutworms Leafhoppers Loopers Melonworms Plant Bugs	4-8 oz/A	0 days	7 days
Specific Directions: *Read the label. *Apply by air or ground equipment. By air, apply in a minimum of 4 gallons of water per acre *By ground, apply in a minimum of 20 gallons of water per acre *Use sufficient water to obtain full coverage of foliage				
Eggplant	Flea Beetles Leafminers Loopers	6 oz/A	3 days	7 days
Specific Directions: *Apply as needed, using sufficient water to obtain uniform coverage *Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre *By ground, apply in a minimum of 10 gallons of water per acre *Use sufficient water to obtain full coverage of foliage				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Garlic	Armyworms Thrips	6-18 oz/A	1 day	10 days
	<p>Specific Directions: *Begin applications when pests appear and repeat as necessary to maintain control *Use higher label rates as onion thrips population increases and avoid rescue situations *Do not apply more than 32 oz. product (0.8 lbs.. a.i.) per acre per season *Apply by air or ground equipment. By air, apply in a minimum of 5 gallons of water per acre *By ground, apply in a minimum of 20 gallons of water per acre *Use sufficient water to obtain full coverage of foliage</p>			
	Cutworms Leafminers Stink Bugs	4-8 oz/A	1 day	10 days
	<p>Specific Directions: *Begin applications when pests appear and repeat as necessary to maintain control *Use higher label rates as onion thrips population increases and avoid rescue situations *Do not apply more than 32 oz. product (0.8 lbs.. a.i.) per acre per season *Apply by air or ground equipment. By air, apply in a minimum of 5 gallons of water per acre *By ground, apply in a minimum of 20 gallons of water per acre *Use sufficient water to obtain full coverage of foliage</p>			
Horseradish	Weevils	6 oz/A	30 days	10 days
	<p>Specific Directions: *Read the label *Do not apply more than 18 oz. product (0.45 lbs.. a.i.) per acre per season *As a spring preplant dip, soak sets for 30 minutes and air-dry before planting *For foliar application, apply with ground equipment in a minimum spray volume of 20 gallons finished spray per acre. Make up to 3 foliar applications as needed to control Weevils adults during oviposition</p>			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Leafy Vegetables: Celery Lettuce (Head, Leaf) Parsley	Aphids Armyworms Cutworms Leafminers	4-8 oz/A	1 day	7 days
	Specific Directions: *Read the label			
	Loopers Leafhoppers	2-8 oz/A	1 day	7 days
	Specific Directions: *Read the label			
Onion	Armyworms Thrips	6-12 oz/A	1 day	7 days
	Specific Directions: *Begin applications when pests appear and repeat as necessary to maintain control *Use higher label rates as onion thrips population increases and avoid rescue situations *Do not apply more than 40 oz. product (1.0 lbs.. a.i.) per acre per season *Apply by air or ground equipment. By air, apply in a minimum of 5 gallons of water per acre *By ground, apply in a minimum of 20 gallons of water per acre *Use sufficient water to obtain full coverage of foliage			
	Cutworms Leafminers Stink Bugs	4-12 oz/A	1 day	7 days
	Specific Directions: *Begin applications when pests appear and repeat as necessary to maintain control *Use higher label rates as onion thrips population increases and avoid rescue situations *Do not apply more than 40 oz. product (1.0 lbs.. a.i.) per acre per season *Apply by air or ground equipment. By air, apply in a minimum of 5 gallons of water per acre *By ground, apply in a minimum of 20 gallons of water per acre *Use sufficient water to obtain full coverage of foliage			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Peppers, Bell	Cutworms Flea Beetles Leafminers Loopers Weevils	4-8 oz/A	3 days	5 days
	Specific Directions: *Do not apply more than 32 oz. product (0.8 lbs.. a.i.) per acre per season *Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre *By ground, apply in a minimum of 10 gallons of water per acre			
Tomato	Armyworms Beetles Cutworms Leafminers Loopers	2-8 oz/A	0 days	7 days
	Specific Directions: *Do not apply more than 24 oz. product (0.6 lbs.. a.i.) per acre per season *Do not apply to cherry tomatoes or other varieties which produce fruit less than one inch in diameter *Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre *By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage			

PURESPRAY GREEN OIL **

Fungicide/Insecticide/Miticide
OMRI Listed

PURESPRAY GREEN OIL **

See label

SAFER INSECTICIDAL SOAP **

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Beans	Aphids Grasshoppers Spider Mites Whiteflies	2.5 fl oz/gal water or 5 Tbsp/gal water	0 days	See label
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Apply when insects first appear or when damage occurs *Spray to wet all infested plant surfaces *One gallon of spray covers approximately 850 sq. ft. of growing area *For most pests repeat application after several days to eliminate newly hatched individuals *Do not make more than 3 sequential applications over a 2 week period *Do not apply during heat of day or when leaf temperature exceeds 90°F 			
Brassica Leafy Vegetables: Broccoli Cabbage Chinese Cabbage	Aphids (Suppression) Leafhoppers Plant Bugs Whiteflies	2.5 fl oz/gal water or 5 Tbsp/gal water	0 days	See label
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Apply when insects first appear or when damage occurs *Spray to wet all infested plant surfaces *One gallon of spray covers approximately 850 sq. ft. of growing area *For most pests repeat application after several days to eliminate newly hatched individuals *Do not make more than 3 sequential applications over a 2 week period. *Do not apply during heat of day or when leaf temperature exceeds 90°F 			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
<p>Carrot Radish</p>	<p>Aphids (Suppression) Leafhoppers Plant Bugs Whiteflies</p>	<p>2.5 fl oz/gal water or 5 Tbsp/gal water</p>	<p>0 days</p>	<p>See label</p>
<p>Specific Directions: *Read the label *Apply when insects first appear or when damage occurs *Spray to wet all infested plant surfaces *One gallon of spray covers approximately 850 sq. ft. of growing area. *For most pests repeat application after several days to eliminate newly hatched individuals *Do not make more than 3 sequential applications over a 2 week period. *Do not apply during heat of day or when leaf temperature exceeds 90°F</p>				
<p>Citrus: Grapefruit Lemon Orange</p>	<p>Aphids Scales Mites</p>	<p>2.5 fl oz/gal water or 5 Tbsp/gal water</p>	<p>0 days</p>	<p>7 days</p>
<p>Specific Directions: *Read the label *Spray when insects first appear or when their populations attain spray level *Apply to achieve thorough coverage of infested branches and/or foliage *Repeat at weekly intervals or as needed *Avoid spraying when blossoms are present *Some marking of fruit may occur, especially when spray run-off occurs or when sooty Mold is present</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Cucurbits: Cantaloupe Cucumber Honeydew Muskmelon Pepino Squash, Summer Watermelon	Aphids (Suppression) Mites Plant Bugs Whiteflies	2.5 fl oz/gal water or 5 Tbsp/gal water	0 days	See label
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Always apply to cucumbers at night or during the coolest part of the day *Spray to wet all infested plant surfaces. *One gallon of spray covers approximately 850 sq. ft. of growing area *Do not make more than 3 sequential applications over a 2 week period *Do not apply during heat of day or when leaf temperature exceeds 90°F 				
Eggplant Pepper (Bell, Hot) Tomato	Aphids (Suppression) Mites Whiteflies	2.5 fl oz/gal water or 5 Tbsp/gal water	0 days	See label
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Apply when insects first appear or when damage occurs *Spray to wet all infested plant surfaces *One gallon of spray covers approximately 850 sq. ft. of growing area *For most pests repeat application after several days to eliminate newly hatched individuals *Do not make more than 3 sequential applications over a 2 week period *Do not apply during heat of day or when leaf temperature exceeds 90°F 				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Garlic	Thrips (Suppression)	2.5 fl oz/gal water or 5 Tbsp/gal water	0 days	See label
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Apply when insects first appear or when damage occurs *Spray to wet all infested plant surfaces *One gallon of spray covers approximately 850 sq. ft. of growing area *For most pests repeat application after several days to eliminate newly hatched individuals *Do not make more than 3 sequential applications over a 2 week period. *Do not apply during heat of day or when leaf temperature exceeds 90°F 			
Lettuce (Head, Leaf)	Aphids (Suppression) Mites Whiteflies	2.5 fl oz/gal water or 5 Tbsp/gal water	0 days	See label
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Apply when insects first appear or when damage occurs *Spray to wet all infested plant surfaces *Do not make more than 3 sequential applications over a 2 week period *Do not apply during heat of day or when leaf temperature exceeds 90°F 			

SEVIN 4F

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Brassica Leafy Vegetables: Broccoli Cauliflower Cabbage Chinese Cabbage Collards Mustard Greens	Flea Beetles Leafhoppers	0.5-1 qts/A	See label	7 days
	<p>Specific Directions: *Read the label *Repeat applications as needed up to a total of 4 times but not more than once every 7 days <u>RESTRICTIONS AND PRECAUTIONS: BRASSICA LEAFY VEGETABLES</u> *For Broccoli and Cabbage Cauliflower do not apply within 3 days of harvest *For Chinese Cabbage, Collards, Kale, and Mustard Greens, do not apply within 14 days of harvest *Do not apply more than a total of 6 quarts per acre per crop</p>			
	Armyworms Caterpillars Diamondback Moths Plant Bugs Stink Bugs	1-2 qts/A	3 See label	7 days
<p>Specific Directions: *Read the label *Repeat applications as needed up to a total of 4 times but not more than once every 7 days <u>RESTRICTIONS AND PRECAUTIONS: BRASSICA LEAFY VEGETABLES</u> *For Broccoli and Cabbage Cauliflower do not apply within 3 days of harvest *For Chinese Cabbage, Collards, Kale, and Mustard Greens, do not apply within 14 days of harvest *Do not apply more than a total of 6 quarts per acre per crop</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Citrus:	Cutworms Leafrollers	2-3 qts/A	5 days	14 days
Grapefruit Lemon Lime Orange Tangerine	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Repeat applications as necessary up to a total of 8 times but not more than once every 14 days *Do not apply more than a total of 20 qts/A/crop *Observe bee caution *Apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, and density, and other factors *For scale control, apply when crawlers are present *For best control of Eriophyid mites including citrus rust mite, apply when pest populations are low 			
	Mites Plant Bugs Scales	3-5 qts/A	5 days	14 days
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Repeat applications as necessary up to a total of 8 times but not more than once every 14 days *Do not apply more than a total of 20 qts/A/crop *Observe bee caution *Apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, and density, and other factors *For scale control, apply when crawlers are present *For best control of Eriophyid mites including citrus rust mite, apply when pest populations are low 			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Beetles Weevils	5-7.5 qts/A	5 days	14 days
<p>Specific Directions: *Read the label *Repeat applications as necessary up to a total of 8 times but not more than once every 14 days *Do not apply more than a total of 20 qts/A/crop *Observe bee caution *Apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, and density, and other factors *For scale control, apply when crawlers are present *For best control of Eriophyid mites including citrus rust mite, apply when pest populations are low</p>				
Corn, Field	Armyworms Flea Beetles Japanese Beetles Leafhoppers	1-2 qt/A	See label	14 days
<p>Specific Directions: *Read the label *Observe bee caution *Repeat applications as needed up to a total of 4 times but not more than once every 14 days *Do not apply more than a total of 8 qts/acre/crop *Optimum timing and good coverage are essential for effective control *Do not apply within 48 days of harvest of grain and fodder or within 14 days of harvest or grazing of forage or silage</p>				
<p>Cutworms</p> <p>2 qt/A</p> <p>See label</p> <p>14 days</p> <p>Specific Directions: *Read the label *Observe bee caution *Repeat applications as needed up to a total of 4 times but not more than once every 14 days *Do not apply more than a total of 8 qts/acre/crop *Optimum timing and good coverage are essential for effective control *Do not apply within 48 days of harvest of grain and fodder or within 14 days of harvest or grazing of forage or silage</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Corn, Sweet	Armyworms Flea Beetles Japanese Beetles Leafhoppers	1-2 qt/A	See label	3 days
	Specific Directions: *Read the label *Observe bee caution *Repeat applications as needed up to a total of 8 times but not more than once every 3 days *Do not apply more than a total of 16 qts/acre/crop *Optimum timing and good coverage are essential for effective control *Do not apply within 2 days of harvest of ears, within 14 days of harvest or grazing of forage, or within 48 days of fodder			
	Cutworms	2 qt/A	See label	3 days
Specific Directions: *Read the label *Observe bee caution *Repeat applications as needed up to a total of 8 times but not more than once every 3 days *Do not apply more than a total of 16 qts/acre/crop *Optimum timing and good coverage are essential for effective control *Do not apply within 2 days of harvest of ears, within 14 days of harvest or grazing of forage, or within 48 days of fodder				
Cucurbits: Cucumbers Melons Pumpkins Squash	Caterpillars Melonworms	0.5-1 qt/A	3 days	7 days
Specific Directions: *Read the label *Repeat applications as necessary up to a total of 6 times but not more often than once every 7 days *Do not apply more than a total of 6 qts/acre/crop *Observe plant response precautions *For optimum control of Squash Bugs, apply sufficient spray volume for thorough coverage and time sprays for early morning or late afternoon				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL				
	Cucumber Beetles Flea Beetles Leafhoppers Squash Bugs	1 qt/A	3 days	7 days				
<p>Specific Directions: *Read the label *Repeat applications as necessary up to a total of 6 times but not more often than once every 7 days *Do not apply more than a total of 6 qts/acre/crop *Observe plant response precautions *For optimum control of Squash Bugs, apply sufficient spray volume for thorough coverage and time sprays for early morning or late afternoon</p>								
<p>Fruiting Vegetables: Eggplant Peppers (Bell, Hot) Tomatoes</p>	Armyworms Beetles Caterpillars Plant Bugs Stink Bugs (Suppression) Thrips (Suppression)	1-2 qt/A	3 days	7 days				
<p>Specific Directions: *Read the label *Repeat applications as necessary up to a total of 7 times but not more often than once every 7 days *Do not apply more than a total of 8 qts/acre/crop *Thorough coverage is essential to effectively suppress stink Bugs</p>								
<table border="1"> <tr> <td data-bbox="412 1457 678 1562">Flea Beetles Leafhoppers</td> <td data-bbox="678 1457 943 1562">0.5-1 qt/A</td> <td data-bbox="943 1457 1208 1562">3 days</td> <td data-bbox="1208 1457 1468 1562">7 days</td> </tr> </table>					Flea Beetles Leafhoppers	0.5-1 qt/A	3 days	7 days
Flea Beetles Leafhoppers	0.5-1 qt/A	3 days	7 days					
<p>Specific Directions: *Read the label *Repeat applications as necessary up to a total of 7 times but not more often than once every 7 days *Do not apply more than a total of 8 qts/acre/crop *Thorough coverage is essential to effectively suppress stink Bugs</p>								

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Cutworms	2 qt/A	3 days	7 days
<p>Specific Directions: *Read the label *Repeat applications as necessary up to a total of 7 times but not more often than once every 7 days *Do not apply more than a total of 8 qts/acre/crop *Thorough coverage is essential to effectively suppress stink Bugs</p>				
<p>Leafy Vegetables: Celery Lettuce (Head, Leaf) Parsley</p>	Flea Beetles Leafhoppers	0.5-1 qt/A	14 days	7 days
	<p>Specific Directions: *Read the label *Repeat applications as necessary up to a total of 5 times but not more often than once every 7 days *Do not apply more than a total of 6 qts/acre/crop</p>			
	Armyworms Caterpillars Plant Bugs Stink Bugs	1-2 qt/A	14 days	7 days
<p>Specific Directions: *Read the label *Repeat applications as necessary up to a total of 5 times but not more often than once every 7 days *Do not apply more than a total of 6 qts/acre/crop</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Legumes: Beans (Fresh, Dried)	Caterpillars Cucumber Beetles Cutworms Japanese Beetles Loopers Plant Bugs	0.5-1 qt/A	See label	7 days
	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Repeat applications as necessary up to a total of 4 times but not more often than once every 7 days *Do not apply more than a total of 6 qts/acre/crop *Do not apply within 14 days of grazing or harvest for forage or within 3 days of harvest of fresh beans or peas or within 21 days of harvest of dried peas, seed, or hay *Observe plant response precautions *For cutworm control, this product is most effective against species which feed on the upper portions of the plant *Use lower rates for light to moderate populations and smaller instars and to provide maximum survival of beneficial insects and spider. Use the higher rates for heavy populations and larger instars 			
	Flea Beetles Leafhoppers Thrips	1 qt/A	See label	7 days
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label *Repeat applications as necessary up to a total of 4 times but not more often than once every 7 days *Do not apply more than a total of 6 qts/acre/crop *Do not apply within 14 days of grazing or harvest for forage or within 3 days of harvest of fresh beans or peas or within 21 days of harvest of dried peas, seed, or hay *Observe plant response precautions *For cutworm control, this product is most effective against species which feed on the upper portions of the plant *Use lower rates for light to moderate populations and smaller instars and to provide maximum survival of beneficial insects and spider. Use the higher rates for heavy populations and larger instars 				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Armyworms Cutworms Stink Bugs	1-1.5 qt/A	See label	7 days
<p>Specific Directions: *Read the label *Repeat applications as necessary up to a total of 4 times but not more often than once every 7 days *Do not apply more than a total of 6 qts/acre/crop *Do not apply within 14 days of grazing or harvest for forage or within 3 days of harvest of fresh beans or peas or within 21 days of harvest of dried peas, seed, or hay *Observe plant response precautions *For cutworm control, this product is most effective against species which feed on the upper portions of the plant *Use lower rates for light to moderate populations and smaller instars and to provide maximum survival of beneficial insects and spider. Use the higher rates for heavy populations and larger instars</p>				
Okra	Stink Bugs	1-1.5 qt/A	3 days	8 days
<p>Specific Directions: *Read the label *Do not apply more than a total of 6 qts/A/crop *Apply using ground equipment in sufficient volume to insure good coverage *Apply as needed on 6 to 8 day intervals</p>				
Root and Tuber Crops: Carrots Horseradish Radish Potato	Flea Beetles Leafhoppers	0.5-1 qt/A	7 days	7 days
<p>Specific Directions: *Repeat applications as necessary up to a total of 6 times but not more often than once every 7 days *Do not apply more than a total of 6 quarts per acre per crop</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Armyworms Beetles Caterpillars Cutworms Plant Bugs Stink Bugs	1-2 qt/A	7 days	7 days
<p>Specific Directions: *Read the label *Repeat applications as necessary up to a total of 6 times but not more often than once every 7 days *Do not apply more than a total of 6 qts/acre/crop *For cutworm control, this product is most effective against species which feed on the upper portions of the plant</p>				
Sweet Potato	Armyworms Beetles Caterpillars Cucumber Beetles Flea Beetles Weevils	1-2 qt/A	7 days	7 days
<p>Specific Directions: *Read the label *For foliar sprays, repeat applications as necessary up to a total of 8 times but not more often than once every 7 days *Do not apply more than a total of 8 qts/acre/crop with in-season sprays *Do not apply more than a total of 1.2 qts/A as a preplant dip treatment</p>				

SEVIN SL

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Brassica Leafy Vegetables: Broccoli Cabbage Chinese Cabbage (Napa)	Flea Beetles Leafhoppers	0.5-1 qts/A	3 days	7 days
	Specific Directions: *Repeat applications as needed up to a total of 4 times *Do not apply more than a total of 6 quarts per acre per crop per year (6 pounds active ingredient per acre per crop per year) *Application is permitted only within 30 days from the date of crop emergence or the date of transplanting *Bee caution: do not apply this product to target crops or weeds in bloom			
	Armyworms Diamondback Moths Stink Bugs Plant Bugs	1-2 qts/A	3 days	7 days
Specific Directions: *Repeat applications as needed up to a total of 4 times *Do not apply more than a total of 6 quarts per acre per crop per year (6 pounds active ingredient per acre per crop per year) *Application is permitted only within 30 days from the date of crop emergence or the date of transplanting *Bee caution: do not apply this product to target crops or weeds in bloom				
Brassica Leafy Vegetables: Chinese Cabbage (Bok Choy) Collards Mustard Greens	Flea Beetles Leafhoppers	0.5-1 qts/A	14 days	7 days
	Specific Directions: *Repeat applications as needed up to a total of 14 times but not more often than once every 7 days *Do not apply more than a total of 6 quarts per acre per crop per year (6 pounds active ingredient per acre per crop per year) *Application is permitted only within 30 days from the date of crop emergence or the date of transplanting *Bee caution: do not apply this product to target crops or weeds in bloom			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Armyworms Diamondback Moths Stink Bugs Plant Bugs	1-2 qts/A	14 days	7 days
<p>Specific Directions:</p> <p>*Repeat applications as needed up to a total of 14 times but not more often than once every 7 days</p> <p>*Do not apply more than a total of 6 quarts per acre per crop per year (6 pounds active ingredient per acre per crop per year)</p> <p>*Application is permitted only within 30 days from the date of crop emergence or the date of transplanting</p> <p>*Bee caution: do not apply this product to target crops or weeds in bloom</p>				
Citrus: Grapefruit Lemon Lime Orange	Cutworms Leafrollers	2-3 qts/A	5 days	14 days
<p>Specific Directions:</p> <p>*Repeat applications as needed up to a total of 8 times but not more often than once every 14 days</p> <p>*Bee caution: do not apply this product to target crops or weeds in bloom. Moving weeds or cover crop bloom prior to applying this product will reduce hazard to bees. Not for use on trees intended for resale of produce or other commercial use. Not for use on trees covering areas greater than 1,000 Ft²</p>				
<p>Mites Plant Bugs Scales</p>				
<p>Specific Directions:</p> <p>*Repeat applications as needed up to a total of 8 times but not more often than once every 14 days</p> <p>*Bee caution: do not apply this product to target crops or weeds in bloom. Moving weeds or cover crop bloom prior to applying this product will reduce hazard to bees. Not for use on trees intended for resale of produce or other commercial use. Not for use on trees covering areas greater than 1,000 Ft²</p> <p>*For scale, apply when pest populations are low</p> <p>*For best results on Eriophyid Mites, including citrus rust mite, apply when pest populations are low</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Weevils	5 qts/A	5 days	14 days
<p>Specific Directions: *Repeat applications as needed up to a total of 8 times but not more often than once every 14 days *Bee caution: do not apply this product to target crops or weeds in bloom. Moving weeds or cover crop bloom prior to applying this product will reduce hazard to bees. Not for use on trees intended for resale of produce or other commercial use. Not for use on trees covering areas greater than 1,000 Ft²</p>				
Fruiting Vegetables: Eggplant Pepinos Pepper (Bell, Chili, Cooking, Sweet) Tomato	Armyworms Stink Bugs (suppression) Thrips (suppression)	1-2 qts/A	3 days	7 days
	<p>Specific Directions: *Repeat applications as needed up to a total of 7 times but not more often than once every 7 days *Thorough coverage is necessary to effectively suppress Stink Bugs *When disease transmission is suspected, monitor fields following application and retreat if reinfestation occurs but not more often than once every 7 days *Do not apply more than a total of 8 quarts per acre per crop per year (8 pounds active ingredient per acre per crop per year) *Bee caution: do not apply this product to target crops or weeds in bloom</p>			
	Flea Beetles Leafhoppers	0.5-1 qts/A	3 days	7 days
<p>Specific Directions: *Repeat applications as needed up to a total of 7 times but not more often than once every 7 days *Thorough coverage is necessary to effectively suppress Stink Bugs *When disease transmission is suspected, monitor fields following application and retreat if reinfestation occurs but not more often than once every 7 days *Do not apply more than a total of 8 quarts per acre per crop per year (8 pounds active ingredient per acre per crop per year) *Bee caution: do not apply this product to target crops or weeds in bloom</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Cutworms	2 qts/A	3 days	7 days
	<p>Specific Directions:</p> <p>*Repeat applications as needed up to a total of 7 times but not more often than once every 7 days</p> <p>*Thorough coverage is necessary to effectively suppress Stink Bugs</p> <p>*When disease transmission is suspected, monitor fields following application and retreat if reinfestation occurs but not more often than once every 7 days</p> <p>*Do not apply more than a total of 8 quarts per acre per crop per year (8 pounds active ingredient per acre per crop per year)</p> <p>*Bee caution: do not apply this product to target crops or weeds in bloom</p> <p>*For cutworm, this product is more effective against species which feed on the upper portions of the plant</p>			
<p>Leafy Vegetables:</p> <p>Celery</p> <p>Lettuce (Head, Leaf)</p> <p>Parsley</p>	Flea Beetles Leafhoppers	0.5-1 qts/A	14 days	7 days
	<p>Specific Directions:</p> <p>*Repeat applications as needed up to a total of 5 times but not more often than once every 7 days</p> <p>*Do not apply more than a total of 6 quarts per acre per crop per year (6 pounds active ingredient per acre per crop per year)</p> <p>*Bee caution: do not apply this product to target crops or weeds in bloom</p>			
	Armyworms Stink Bugs Plant Bugs	1-2 qts/A	14 days	7 days
	<p>Specific Directions:</p> <p>*Repeat applications as needed up to a total of 5 times but not more often than once every 7 days</p> <p>*Do not apply more than a total of 6 quarts per acre per crop per year (6 pounds active ingredient per acre per crop per year)</p> <p>*Bee caution: do not apply this product to target crops or weeds in bloom</p>			

Sluggo SNAIL & SLUG BAIT **

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Avocado Grapefruit Lemon Lime Orange Tangerine	Slugs Snails	40 lbs/A	0	As needed
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label. *Broadcast liberally under trees, around irrigation pipes, farm buildings and any other place where slugs and snails may congregate. *Use only as a baiting treatment to be applied directly to the soil area beneath plants. *Do not allow bait pellet to come in contact with parts of plant that may be used for food or feed purposes. *Slug & Snail Bait should not be applied to dry soil. *Applications should be made following an irrigation for best results. *Apply any time of the day, but evening applications are preferred. *Do not re-water for 48 hours after application. 				
Beans Bittermelon Broccoli	Slugs Snails	30-40 lbs/A or 1 lb/1,100 sq Ft	0	As needed
<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label. *Use only as a baiting treatment to be applied directly to the soil area beneath plants. *Do not allow bait pellet to come in contact with parts of plant that may be used for food or feed purposes. *Slug & Snail Bait should not be applied to dry soil. *Applications should be made following an irrigation for best results. *Apply any time of the day, but evening applications are preferred. *Do not re-water for 48 hours after application. 				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Cabbage Cantaloupe Carrot Cauliflower	Slugs Snails	1,500 sq. ft. of soil surface	0	As needed
Corn Cucumber Eggplant Garlic Gourds Honeydew Horseradish Lettuce Muskmelon Mustard Greens Okra Pepino Pepper, Bell Pepper, Hot Pumpkin Radish Squash, Summer Squash, Winter Sweet Potato Tomato Watermelon	<p>Specific Directions:</p> <ul style="list-style-type: none"> *Read the label. *Use only as a baiting treatment to be applied directly to the soil area beneath plants. *Do not allow bait pellet to come in contact with parts of plant that may be used for food or feed purposes. *Slug & Snail Bait should not be applied to dry soil. *Applications should be made following an irrigation for best results. *Apply any time of the day, but evening applications are preferred. *Do not re-water for 48 hours after application. 			

TRIACT 70 **

Fungicide/Insecticide/Miticide
OMRI Listed

TRIACT 70 **

See label

VENDEX 50WP

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Citrus: Grapefruit Lemon Lime Orange Tangerine	Mites	2-4 lbs./A	7 days	See label
	Specific Directions: *Read the label *Apply when Mites first appear *Make no more than 2 applications per year			
Eggplant	Mites	2-3 lbs./A	3 days	See label
	Specific Directions: *Read the label *Apply when Mites first appear *Repeat as necessary to maintain mite control, making no more than 3 applications per year *Apply no more than 9 lbs. per acre per year			
Papaya	Mites	2 lbs./A	7 days	See label
	Specific Directions: *Read the label *Apply when Mites first appear *Repeat as necessary to maintain mite control, making no more than 4 applications per season *Apply no more than 8 lbs. per acre per year *Do not apply more than once per month to bearing trees			

VYDATE L

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Cantaloupe Cucumber Honeydew Melon Pumpkin Squash (Summer, Winter) Watermelon	Aphids Leafminers	2-4 pt/A	1 day	7 days
	Specific Directions: *Read the label *Do not apply more than 16 pts (2 gals) Vydate L per acre per season *Apply in sufficient water to obtain uniform coverage of foliage *Make applications when insects first appear *Apply the low rate for light infestations of insects and the high rate for severe infestations *Where Leafminer infestations occur annually, initiate treatment schedule 2-4 weeks after planting and repeat weekly or as needed			
	Nematodes	See label	See label	See label
Carrot	Specific Directions: *Read the label *Do not apply more than 16 pts (2 gals) Vydate L per acre per season *Apply in sufficient water to obtain uniform coverage of foliage *Make applications when insects first appear *Apply the low rate for light infestations of insects and the high rate for severe infestations *Where Leafminer infestations occur annually, initiate treatment schedule 2-4 weeks after planting and repeat weekly or as needed			
	Nematodes	See label	See label	See label
Specific Directions: *Read the label *Do not apply more than 20 pts (2.5 gals) Vydate L per acre per season				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Citrus (Fruit-bearing)	Mites	0.25-1 pt/100 gal water	7 days	See label
Grapefruit Lemon Lime Orange Tangerine	<p>Specific Directions: *Read the label *Spray to run-off using up to 400 gals of water per acre *Do not apply more than 4 qts of product per acre *Begin applications as soon as significant infestations are found and continue on an as needed basis *Under light to moderate pressure, apply on 4 to 6 week intervals or as needed; whereas under moderate to heavy pressure, apply every 2 to 3 weeks as long as the infestation continues *Do not make more than 6 applications/year NOTE: Do not graze livestock in treated orchards. Do not handle plants or come in contact with foliage within 3 days of application. This product is toxic to bees and should not be applied when bees are actively visiting the area *Do not apply more than 28 pts (3.5 gals) Vydate L per acre per year</p>			
	Thrips	2-4 pt/A	7 days	See label
	<p>Specific Directions: *Read the label *Apply in sufficient water (100-500 gals per acre) to give good coverage NOTE: Do not graze livestock in treated orchards. Do not handle plants or come in contact with foliage within 3 days of application. This product is toxic to bees and should not be applied when bees are actively visiting the area *Do not apply more than 28 pts (3.5 gals) Vydate L per acre per year</p>			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
<p>Citrus (Non Fruit-bearing)</p> <p>Grapefruit Lemon Lime Orange Tangerine</p>	<p>Aphids Beetles Leafhoppers Leafminers Mealybugs Mites Scales Thrips Whiteflies (Foliar Treatment)</p>	<p>See label</p>	<p>See label</p>	<p>See label</p>
<p>Specific Directions: *Read the label NOTE: Use only on commercial plantings; do not use on home plantings. Since varieties are numerous, continually changing, and may respond differently to Vydate L, test Vydate L on a small scale before proceeding to large-scale application. Varietal response may also vary if Vydate L is mixed with other products *Do not apply more than 32 pts (4 gals) per acre per season</p>				
<p>Nematodes</p>		<p>See label</p>	<p>See label</p>	<p>See label</p>
<p>Specific Directions: *Read the label NOTE: Use only on commercial plantings; do not use on home plantings. Since varieties are numerous, continually changing, and may respond differently to Vydate L, test Vydate L on a small scale before proceeding to large-scale application. Varietal response may also vary if Vydate L is mixed with other products *Do not apply more than 32 pts (4 gals) per acre per season</p>				
<p>Eggplant</p>	<p>Aphids Beetles Leafminers Mites</p>	<p>2-4 pts/A</p>	<p>1 day</p>	<p>10 days-3 weeks</p>
<p>Specific Directions: *Read the label. *Do not apply more than 12 pts (1.5 gals) Vydate L per acre per season</p>				

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
	Nematodes	See label	See label	See label
	<p>Specific Directions: *Read the label. *Do not apply more than 24 pts (3 gals) Vydate L per acre per season</p>			
Pepper (Bell, Hot)	Aphids Leafminers Weevils (Foliar Treatment)	2-4 pt/A	7 days	1-2 weeks
	<p>Specific Directions: *Read the label *Do not apply more than 14 pts (1.75 gals) Vydate L per acre per season *Apply in sufficient water to obtain uniform coverage *Make applications when insects first appear *Apply low rates for light infestations of insects *Use highest recommended rates and shorter intervals for controlling severe infestations</p>			
	Nematodes	See label	See label	See label
	<p>Specific Directions: *Read the label *Do not apply more than 14 pts (1.75 gals) Vydate L per acre per season *Apply in sufficient water to obtain uniform coverage *Make applications when insects first appear *Apply low rates for light infestations of insects *Use highest recommended rates and shorter intervals for controlling severe infestations</p>			
Pineapple	Nematodes	See label	See label	See label
	<p>Specific Directions: *Read the label *Do not apply more than 4 gals Vydate L per acre per year *Do not harvest or graze treated fields within 30 days of application</p>			

CROP	PEST(S) CONTROLLED	RATE	DAYS TO HARVEST	SPRAYING INTERVAL
Sweet Potato	Nematodes	See label	See label	See label
	Specific Directions: *Read the label *Do not apply more than 24 pts (3 gals) Vydate L per acre per season			
Tomato	Aphids Beetles Leafminers	2-4 pt/A	3 days	5-7 days
	Specific Directions: *Read the label *Do not apply more than 32 pts (4 gals) Vydate L per acre per season *Apply in sufficient water to obtain uniform coverage *Make applications when insects first appear *Apply low rates for light infestations of insects *Use highest recommended rates and shorter intervals for controlling severe infestations			
	Nematodes	See label	See label	See label
	Specific Directions: *Read the label *Do not apply more than 32 pts (4 gals) Vydate L per acre per season *Apply in sufficient water to obtain uniform coverage *Make applications when insects first appear *Apply low rates for light infestations of insects *Use highest recommended rates and shorter intervals for controlling severe infestations			

Crop and Pest Use Index

Avocado

Insects, Mites and Nematodes

Ants

Amdro.....p.22

Aphids

Alias 4Fp.21

Caterpillars

Dipel DF.....p.36

Perm-Up 3.2 EC.....p.95

Leafhoppers

Alias 4Fp.21

Perm-Up 3.2 EC.....p.95

Leafrollers

Dipel DF.....p.36

Lannate LV.....p.67

Loopers

Dipel DF.....p.36

Lannate LV.....p.67

Malathion 5 EC.....p.76

Malathion 8

AQUAMUL.....p.82

Perm-Up 3.2 EC.....p.95

Mealybugs

Alias 4Fp.21

Scales

Alias 4Fp.21

Malathion 5 EC.....p.76

Malathion 8

AQUAMUL.....p.82

Perm-Up 3.2 EC.....p.95

Scale Crawlers

Perm-Up 3.2 EC.....p.95

Thrips

Alias 4Fp.21

Malathion 5 EC.....p.76

Malathion 8

AQUAMUL.....p.82

Perm-Up 3.2 EC.....p.95

Whiteflies

Alias 4Fp.21

Perm-Up 3.2 EC.....p.95

Diseases

Anthracnose

Kocide 3000p.59

Blotch

Kocide 3000p.59

Scab

Kocide 3000p.59

Other

CROP AND PEST USE INDEX

Slugs and Snails

Sluggo Snail & Slug Bait.....p.119

Banana

Insects, Mites and Nematodes

Ants

Amdro Fire Ant Bait.....p.22

Aphids

Alias 4Fp.17

Banana Skipper

Dipel DF.....p.36

Leafhoppers

Alias 4Fp.17

Thrips

Alias 4Fp.17

Diseases

Sigatoka

Kocide 3000.....p.59

Manzate Pro Stick.....p.88

Beans

Insects, Mites and Nematodes

Aphids

Alias 4F	p.19
Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.33
D.Z.N. Diazinon.....	p.40
Lannate LV.....	p.67
Safer Insecticidal Soap.....	p.102

Armyworms

Dipel DF.....	p.36
Lannate LV.....	p.67

Beetles

Dimethoate 4 EC.....	p.33
D.Z.N. Diazinon.....	p.40
Lannate LV.....	p.67

Caterpillars

Dipel DF.....	p.36
Lannate LV.....	p.67
Sevin 4F.....	p.112

Cucumber Beetles

D.Z.N. Diazinon.....	p.40
Sevin 4F.....	p.112

Cutworms

D.Z.N. Diazinon.....	p.40
Lannate LV.....	p.67
Sevin 4F.....	p.112

Grasshoppers

Dimethoate 4 EC.....	p.32
Safer Insecticidal Soap.....	p.102

Japanese Beetles

Sevin 4F.....	p.112
---------------	-------

Leafhoppers

Alias 4F	p.19
Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.33
Lannate LV.....	p.67

Leafminers

Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.33
D.Z.N. Diazinon.....	p.40

Loopers

Dibrom 8 Emulsive.....	p.30
Dipel DF.....	p.36
Lannate LV.....	p.67
Sevin 4F.....	p.112

Mites

Dimethoate 4 EC.....	p.33
----------------------	------

Plant Bugs

Dimethoate 4 EC.....	p.33
Sevin 4F.....	p.112

Spider Mites

Dibrom 8 Emulsive.....	p.30
Safer Insecticidal Soap.....	p.102

Stink Bugs

Dibrom 8 Emulsive.....	p.30
------------------------	------

Thrips

Lannate LV.....	p.67
-----------------	------

Whiteflies

Alias 4F	p.19
Safer Insecticidal Soap.....	p.102

Diseases

Anthracnose

Bravo 720p.23
Equis 720.....p.50

Ascochyta Blight

Equis 720.....p.50

Brown Spot

Kocide 3000.....p.59

Cercospora Leaf Spot

Bravo 720p.23
Equis 720.....p.50

Common Blight

Kocide 3000.....p.59

Downy Mildew

Equis 720.....p.50

Botrytis Blight (Gray Mold)

Bravo 720p.23
Equis 720.....p.50

Rust

Bravo 720p.23
Equis 720.....p.50

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.119

Betel Nut

Consult your agricultural extension agent for information on pesticide use for this crop.

Bitter Melon

Insects, Mites and Nematodes

Aphids

- Alias 4F.....p.18
- Lannate LV.....p.74
- Perm-Up 3.2 EC.....p.97

Armyworms

- Dipel DF.....p.36
- Lannate LV.....p.74

Caterpillars

- Lannate LV.....p.74

Cucumber Beetles

- Alias 4F.....p.18
- Lannate LV.....p.74
- Perm-Up 3.2 EC.....p.97

Cutworms

- Lannate LV.....p.74
- Perm-Up 3.2 EC.....p.97

Flea Beetles

- Lannate LV.....p.74

Leafhoppers

- Alias 4F.....p.18
- Perm-Up 3.2 EC.....p.97

Leafminers

- Perm-Up 3.2 EC.....p.97

Loopers

- Dipel DF.....p.36
- Lannate LV.....p.74
- Perm-Up 3.2 EC.....p.97

Melonworms

- Dipel DF.....p.36
- Lannate LV.....p.74
- Perm-Up 3.2 EC.....p.97

Plant Bugs

- Perm-Up 3.2 EC.....p.97

Squash Bugs

- Perm-Up 3.2 EC.....p.97

Thrips

- Alias 4F.....p.18

Whiteflies

- Alias 4F.....p.18

Diseases

Anthracnose

- Equis 720.....p.52

Downy Mildew

- Equis 720.....p.52

Target Spot

- Equis 720.....p.52

Other

Slugs and Snails

- Sluggo Snail & Slug Bait.....p.119

Broccoli

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.17
Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.33
D.Z.N. Diazinon.....	p.40
Malathion 5 EC.....	p.76
Malathion 8	

AQUAMUL.....p.82

Perm-Up 3.2 EC.....	p.95
Safer Insecticidal Soap.....	p.102

Armyworms

Dipel DF.....	p.39
Perm-Up 3.2 EC.....	p.95
Sevin 4F.....	p.106
Sevin SL.....	p.115

Cabbageworms

Dibrom 8 Emulsive.....	p.30
------------------------	------

Caterpillars

Dipel DF.....	p.39
D.Z.N. Diazinon.....	p.40
Lannate LV.....	p.68
Malathion 5 EC.....	p.76
Malathion 8	

AQUAMUL.....p.82

Sevin 4F.....	p.106
---------------	-------

Cutworms

Dipel DF.....	p.39
D.Z.N. Diazinon.....	p.40

Diamondback Moths

Dipel DF.....	p.39
D.Z.N. Diazinon.....	p.40
Lannate LV.....	p.68
Perm-Up 3.2 EC.....	p.95
Sevin 4F.....	p.106

Sevin SL.....p.115

Diamondback Caterpillars

Dibrom 8 Emulsive.....p.30

Flea Beetles

Alias 4F.....	p.17
Sevin 4F.....	p.106
Sevin SL.....	p.115

Leafhoppers

Alias 4F.....	p.17
Safer Insecticidal Soap.....	p.102
Sevin 4F.....	p.106
Sevin SL.....	p.115

Leafminers

D.Z.N. Diazinon.....p.40

Leafrollers

Dipel DF.....p.39

Loopers

Dipel DF.....	p.39
Lannate LV.....	p.68
Malathion 5 EC.....	p.76
Malathion 8	
AQUAMUL.....	p.82
Perm-Up 3.2 EC.....	p.95

Plant Bugs

Perm-Up 3.2 EC.....	p.95
Safer Insecticidal Soap.....	p.102
Sevin 4F.....	p.106
Sevin SL.....	p.115

Stink Bugs

Sevin 4F.....	p.106
Sevin SL.....	p.115

Thrips

Perm-Up 3.2 EC.....p.95

CROP AND PEST USE INDEX

Whiteflies

Alias 4F.....p.17

Safer Insecticidal Soap.....p.102

Diseases

Alternaria Leaf Spot

Bravo 720.....p.23

Equis 720.....p.51

Kocide 3000.....p.62

Manzate Pro Stick.....p.88

Downy Mildew

Equis 720.....p.51

Kocide 3000.....p.62

Manzate Pro Stick.....p.88

Black Rot (*Xanthomonas*)

Kocide 3000.....p.62

Other

Slugs and Snails

Deadline GT.....p.48

Sluggo Snail & Slug Bait.....p.119

Cabbage

Insects, Mites and Nematodes

Aphids

Alias 4F.....p.17
 Dibrom 8 Emulsive.....p.30
 Dimethoate 4 EC.....p.33
 D.Z.N. Diazinon.....p.41/43
 Malathion 5 EC.....p.76
 Malathion 8

AQUAMUL.....p.82

Perm-Up 3.2 EC.....p.96
 Safer Insecticidal Soap.....p.102

Armyworms

Dipel DF.....p.39
 Lannate LV.....p.68/70
 Perm-Up 3.2 EC.....p.96
 Sevin 4F.....p.106
 Sevin SL.....p.115/116

Cabbageworms

Dibrom 8 Emulsive.....p.30

Caterpillars

Dipel DF.....p.39
 D.Z.N. Diazinon.....p.41/43
 Lannate LV.....p.68
 Malathion 5 EC.....p.76
 Malathion 8

AQUAMUL.....p.82

Sevin 4F.....p.106

Cutworms

Dipel DF.....p.39
 D.Z.N. Diazinon.....p.41
 Lannate LV.....p.68
 Perm-Up 3.2 EC.....p.96

Diamondback Moths

Dipel DF.....p.39
 D.Z.N. Diazinon.....p.41/43

Lannate LV.....p.68

Malathion 5 EC.....p.76

Malathion 8

AQUAMUL.....p.82

Perm-Up 3.2 EC.....p.96

Sevin 4F.....p.106

Sevin SL.....p.115/116

Diamondback caterpillars

Dibrom 8 Emulsive.....p.30

Flea Beetles

Alias 4F.....p.17

Perm-Up 3.2 EC.....p.96

Sevin 4F.....p.106

Sevin SL.....p.115

Leafhoppers

Alias 4F.....p.17

Safer Insecticidal Soap.....p.102

Sevin 4F.....p.106

Sevin SL.....p.115

Leafminers

D.Z.N. Diazinon.....p.41/43

Leafrollers

Dipel DF.....p.39

Loopers

Dipel DF.....p.39

Lannate LV.....p.68/70

Malathion 5 EC.....p.76

Malathion 8

AQUAMUL.....p.82

Perm-Up 3.2 EC.....p.96

Plant Bugs

Safer Insecticidal Soap.....p.102

Sevin 4F.....p.106

Sevin SL.....p.115/116

CROP AND PEST USE INDEX

Stink Bugs

Sevin 4F.....p.106
Sevin SL.....p.115/116

Alias 4F.....p.17
Safer Insecticidal Soap.....p.102

Whiteflies

Diseases

Alternaria Leaf Spot

Bravo 720.....p.23
Equis 720.....p.51
Kocide 3000.....p.62
Manzate Pro Stick.....p.88

Downy Mildew

Equis 720.....p.51
Kocide 3000.....p.62
Manzate Pro Stick.....p.88

Black Rot (*Xanthomonas*)

Kocide 3000.....p.62

Other

Slugs and Snails

Deadline GT.....p.48
Sluggo Snail & Slug Bait.....p.120

Cacao

Diseases

Black pod

Kocide.....p.59

Cantaloupe

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.34
D.Z.N. Diazinon.....	p.41
Lannate LV.....	p.73
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97
Safer Insecticidal Soap.....	p.104
Vydate L.....	p.123

Armyworms

Dibrom 8 Emulsive.....	p.30
Dipel DF.....	p.36
Lannate LV.....	p.73

Caterpillars

Lannate LV.....	p.73
-----------------	------

Cucumber Beetles

D.Z.N. Diazinon.....	p.41
Lannate LV.....	p.73
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97

Cutworms

D.Z.N. Diazinon.....	p.41
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.97

Flea Beetles

Lannate LV.....	p.73
-----------------	------

Leafhoppers

Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.34
D.Z.N. Diazinon.....	p.41
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97

Leafminers

Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.34
D.Z.N. Diazinon.....	p.41
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97
Vydate L.....	p.123

Loopers

Dibrom 8 Emulsive.....	p.30
Dipel DF.....	p.36
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.97

Melonworms

Dipel DF.....	p.36
D.Z.N. Diazinon.....	p.41
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.97

Mites

Safer Insecticidal Soap.....	p.104
------------------------------	-------

Nematodes

Vydate L.....	p.123
---------------	-------

Plant Bugs

Perm-Up 3.2 EC.....	p.97
Safer Insecticidal Soap.....	p.104

Spider Mites

D.Z.N. Diazinon.....	p.41
Malathion 5 EC.....	p.78

Squash Bugs

Perm-Up 3.2 EC.....	p.97
---------------------	------

Thrips

D.Z.N. Diazinon.....	p.41
----------------------	------

Whiteflies

Safer Insecticidal Soap.....	p.104
------------------------------	-------

Diseases

Anthracnose

Bravo 720.....	p.24
Equis 720.....	p.52
Kocide 3000.....	p.63
Manzate Pro Stick.....	p.91

Alternaria Leaf Blight

Equis 720.....	p.52
----------------	------

Alternaria Leaf Spot

Equis 720.....	p.52
Kocide 3000.....	p.63
Manzate Pro Stick.....	p.91

Angular Leaf Spot

Kocide 3000.....	p.63
------------------	------

Cercospora Leaf Spot

Equis 720.....	p.52
Manzate Pro Stick.....	p.91

Downy Mildew

Equis 720.....	p.52
Kocide 3000.....	p.63
Manzate Pro Stick.....	p.91

Gummy Stem Blight

Equis 720.....	p.52
Kocide 3000.....	p.63
Manzate Pro Stick.....	p.91

Powdery Mildew

Equis 720.....	p.52
Kocide 3000.....	p.63

Scab

Equis 720.....	p.52
----------------	------

Target Spot

Bravo 720.....	p.24
Equis 720.....	p.52

Watermelon Bacterial Fruit Blotch

Kocide 3000.....	p.63
------------------	------

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....	p.120
-------------------------------	-------

Carrot

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.20
D.Z.N Diazinon.....	p.42
Malathion 5 EC.....	p.76
Malathion 8	
AQUAMUL.....	p.82
Safer Insecticidal Soap.....	p.103

Armyworms

Dipel DF.....	p.37
Lannate LV.....	p.69
Sevin 4F.....	p.113

Beetles

Sevin 4F.....	p.113
---------------	-------

Caterpillars

Dipel DF.....	p.37
Sevin 4F.....	p.113

Cutworms

Dipel DF.....	p.37
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.69
Sevin 4F.....	p.113

Diamondback Moths

Dipel DF.....	p.37
---------------	------

Flea Beetles

Alias 4F.....	p.20
Sevin 4F.....	p.113

Leafhoppers

Alias 4F.....	p.20
Lannate LV.....	p.69
Malathion 5 EC.....	p.76
Malathion 8	
AQUAMUL.....	p.82
Safer Insecticidal Soap.....	p.103
Sevin 4F.....	p.113

Leafrollers

Dipel DF.....	p.37
---------------	------

Loopers

Dipel DF.....	p.37
---------------	------

Nematodes

Vydate L.....	p.123
---------------	-------

Plant Bugs

Safer Insecticidal Soap.....	p.103
Sevin 4F.....	p.113

Stink Bugs

Sevin 4F.....	p.113
---------------	-------

Whiteflies

Alias 4F.....	p.20
Safer Insecticidal Soap.....	p.103

Diseases

Cercospora Leaf Spot

Equis 720.....	p.51
Kocide 3000.....	p.60

Early Blight (Cercospora)

Bravo 720.....	p.24
----------------	------

Late Blight (Alternaria)

Bravo 720.....	p.24
Equis 720.....	p.51

CROP AND PEST USE INDEX

Kocide 3000.....p.60

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Cashew

Consult your agricultural extension agent for information on pesticide use for this crop.

Cauliflower

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.17
Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.33
D.Z.N Diazinon.....	p.42
Malathion 5 EC.....	p.77
Perm-Up 3.2 EC.....	p.95

Armyworms

Dipel DF.....	p.39
Perm-Up 3.2 EC.....	p.95
Sevin 4F.....	p.106

Cabbageworms

Dibrom 8 Emulsive.....	p.30
------------------------	------

Caterpillars

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.69
Sevin 4F.....	p.106

Cutworms

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.69

Diamondback Moths

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.69
Malathion 5 EC.....	p.77
Perm-Up 3.2 EC.....	p.95
Sevin 4F.....	p.106

Diamondback Caterpillars

Dibrom 8 Emulsive.....	p.30
------------------------	------

Flea Beetles

Alias 4F.....	p.17
Sevin 4F.....	p.106

Leafhoppers

Alias 4F.....	p.17
Sevin 4F.....	p.106

Leafminers

D.Z.N Diazinon.....	p.42
---------------------	------

Leafrollers

Dipel DF.....	p.39
---------------	------

Loopers

Dipel DF.....	p.39
Lannate LV.....	p.69
Perm-Up 3.2 EC.....	p.95

Plant Bugs

Perm-Up 3.2 EC.....	p.95
Sevin 4F.....	p.106

Stink Bugs

Sevin 4F.....	p.106
---------------	-------

Thrips

Perm-Up 3.2 EC.....	p.95
---------------------	------

Whiteflies

Alias 4F.....	p.17
---------------	------

Diseases

Alternaria Leaf Spot

Bravo 720.....p.23
Equis 720.....p.51

Downy Mildew

Equis 720.....p.51
Kocide 3000.....p.62

Black Leaf Spot

Kocide 3000.....p.62

Black Rot (Xanthomonas)

Kocide 3000.....p.62

Other

Slugs and Snails

Deadline GT.....p.48
Sluggo Snail & Slug Bait.....p.120

Chinese Cabbage

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.17
Dibrom 8 Emulsive.....	p.30
D.Z.N Diazinon.....	p.43
Perm-Up 3.2 EC.....	p. 96
Safer Insecticidal Soap.....	p.102

Armyworms

Dipel DF.....	p.39
Lannate LV.....	p.70
Perm-Up 3.2 EC.....	p. 96
Sevin 4F.....	p.106
Sevin SL.....	p.115/116

Cabbageworm

Dibrom 8 Emulsive.....	p.30
------------------------	------

Caterpillars

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.43
Sevin 4F.....	p.106

Cutworms

Dipel DF.....	p.39
Perm-Up 3.2 EC.....	p. 96

Diamondback Moths

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.43
Perm-Up 3.2 EC.....	p. 96
Sevin 4F.....	p.106
Sevin SL.....	p.115/116

Diamondback Caterpillar

Dibrom 8 Emulsive.....	p.30
------------------------	------

Flea Beetles

Alias 4F.....	p.17
Perm-Up 3.2 EC.....	p. 96
Sevin 4F.....	p.106
Sevin SL.....	p.115

Leafhoppers

Alias 4F.....	p.17
Safer Insecticidal Soap.....	p.102
Sevin 4F.....	p.106
Sevin SL.....	p.115

Leafminers

D.Z.N Diazinon.....	p.43
---------------------	------

Leafrollers

Dipel DF.....	p.39
---------------	------

Loopers

Dipel DF.....	p.39
Lannate LV.....	p.70
Perm-Up 3.2 EC.....	p. 96

Plant Bugs

Safer Insecticidal Soap.....	p.102
Sevin 4F.....	p.106
Sevin SL.....	p.115/116

Stink Bugs

Sevin 4F.....	p.106
Sevin SL.....	p.115/116

Whiteflies

Alias 4F.....	p.17
Safer Insecticidal Soap.....	p.102

Diseases

Alternaria Leaf Spot

- Bravo 720.....p.23
- Equis 720.....p.51
- Kocide 3000.....p.62
- Manzate Pro Stick.....p.90

Anthracnose

- Manzate Pro Stick.....p.90

Black Rot (Xanthomonas)

- Equis 720.....p.51
- Kocide 3000.....p.62

Cercospora Leaf Spot

- Manzate Pro Stick.....p.90

Downy Mildew

- Equis 720.....p.51
- Kocide 3000.....p.62
- Manzate Pro Stick.....p.90

Gummy Stem Blight

- Manzate Pro Stick.....p.90

Scab

- Manzate Pro Stick.....p.90

Corn, Field

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.21
Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.46
Lannate LV.....	p.70

Armyworms

Dipel DF.....	p.37
Lannate LV.....	p.70
Perm-Up 3.2 EC.....	p.96
Sevin 4F.....	p.108/109

Beetles

Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.46
Lannate LV.....	p.70
Malathion 5 EC.....	p.77
Malathion 8	

AQUAMUL.....	p.83
--------------	------

Caterpillars

Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.46
Lannate LV.....	p.70

Cucumber Beetles

Alias 4F.....	p.21
---------------	------

Cutworms

D.Z.N Diazinon.....	p.46
Lannate LV.....	p.71

Sevin 4F.....	p.108/109
---------------	-----------

Flea Beetles

D.Z.N Diazinon.....	p.46
Perm-Up 3.2 EC.....	p.96
Sevin 4F.....	p.108/109

Grasshoppers

Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.46
Lannate LV.....	p.70

Japanese Beetles

Sevin 4F.....	p.108/109
---------------	-----------

Leafhoppers

Alias 4F.....	p.21
Perm-Up 3.2 EC.....	p.96
Sevin 4F.....	p.108/109

Mites

Dimethoate 4 EC.....	p.34
----------------------	------

Spider Mites

Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.46

Whiteflies

Alias 4F.....	p.21
---------------	------

Diseases

Bacterial Stalk Rot

Kocide 3000.....p.62

Helminthosporium Leaf Blight

Bravo 720.....p.24

Equis 720.....p.54

Manzate Pro Stick.....p.89

Rust

Bravo 720.....p.24

Equis 720.....p.54

Manzate Pro Stick.....p.89

Gray Leaf Spot

Manzate Pro Stick.....p.89

Other

Slugs and Snails

Deadline GT.....p.48

Sluggo Snail & Slug Bait.....p.120

Cucumber

Insects, Mites and Nematodes

Aphids

D.Z.N Diazinon.....p.43
Lannate LV.....p.71
Malathion 5 EC.....p.77
Malathion 8

AQUAMUL.....p.83

Perm-Up 3.2 EC.....p.97
Safer Insecticidal Soap.....p.104
Vydate L.....p.123

Armyworms

Dipel DF.....p.36
Lannate LV.....p.71

Beetles

Malathion 5 EC.....p.77
Malathion 8

AQUAMUL.....p.83

Caterpillars

D.Z.N Diazinon.....p.43
Lannate LV.....p.71
Malathion 5 EC.....p.77
Malathion 8

AQUAMUL.....p.83

Sevin 4F.....p.109

Cucumber Beetles

Lannate LV.....p.71
Malathion 5 EC.....p.77
Malathion 8

AQUAMUL.....p.83

Perm-Up 3.2 EC.....p.97
Sevin 4F.....p.110

Cutworms

D.Z.N Diazinon.....p.43
Lannate LV.....p.71
Perm-Up 3.2 EC.....p.97

Diamondback Moths

D.Z.N Diazinon.....p.43

Flea Beetles

Lannate LV.....p.71
Sevin 4F.....p.110

Leafhoppers

Malathion 8

AQUAMUL.....p.83

Perm-Up 3.2 EC.....p.97
Sevin 4F.....p.110

Leafminers

D.Z.N Diazinon.....p.43
Malathion 5 EC.....p.77
Perm-Up 3.2 EC.....p.97
Vydate L.....p.123

Loopers

Dipel DF.....p.36
Perm-Up 3.2 EC.....p.97

Melonworms

Dipel DF.....p.36
Lannate LV.....p.71
Perm-Up 3.2 EC.....p.97
Sevin 4F.....p.109

Mites

Safer Insecticidal Soap.....p.104

Nematodes

Vydate L.....p.123

Plant Bugs

Perm-Up 3.2 EC.....p.97
Safer Insecticidal Soap.....p.104

Spider Mites

CROP AND PEST USE INDEX

Malathion 5 EC.....p.77
Malathion 8
AQUAMUL.....p.83

Squash Bugs

Perm-Up 3.2 EC.....p.97

Sevin 4F.....p.110

Whiteflies

Safer Insecticidal Soap.....p.104

Diseases

Alternaria Leaf Blight

Equis 720.....p.53

Alternaria Leaf Spot

Equis 720.....p.53

Kocide 3000.....p.63

Angular Leaf Spot

Kocide 3000.....p.63

Anthracnose

Bravo 720.....p.24

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.89

Cercospora Leaf Spot

Equis 720.....p.53

Manzate Pro Stick.....p.89

Downy Mildew

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.89

Gummy Stem Blight

Equis 720.....p.53

Kocide 3000.....p.63

Manzate Pro Stick.....p.89

Powdery Mildew

Equis 720.....p.53

Kocide 3000.....p.63

Scab

Equis 720.....p.53

Manzate Pro Stick.....p.89

Target Spot

Bravo 720.....p.24

Equis 720.....p.52

Watermelon Bacterial Fruit Blotch

Kocide 3000.....p.63

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Eggplant

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.19
Dibrom 8 Emulsive.....	p.31
Lannate LV.....	p.71
Malathion 5 EC.....	p.77
Malathion 8	
AQUAMUL.....	p.83
Safer Insecticidal Soap.....	p.104
Vydate L.....	p.125

Armyworms

Dipel DF.....	p.38
Lannate LV.....	p.71
Sevin 4F.....	p.110
Sevin SL.....	p.117

Beetles

Dibrom 8 Emulsive.....	p.31
Sevin 4F.....	p.110
Vydate L.....	p.125

Caterpillars

Dipel DF.....	p.38
Lannate LV.....	p.71
Sevin 4F.....	p.110

Colorado Potato Beetles

Alias 4F.....	p.19
---------------	------

Cutworms

Dipel DF.....	p.38
Sevin 4F.....	p.110
Sevin SL.....	p.118

Flea Beetles

Dibrom 8 Emulsive.....	p.31
Perm-Up 3.2 EC.....	p.97
Sevin 4F.....	p.110
Sevin SL.....	p.117

Leafhoppers

Alias 4F.....	p.19
Sevin 4F.....	p.110
Sevin SL.....	p.117

Leafminers

Dibrom 8 Emulsive.....	p.31
Perm-Up 3.2 EC.....	p.97
Vydate L.....	p.125

Loopers

Dipel DF.....	p.38
Perm-Up 3.2 EC.....	p.97

Mites

Safer Insecticidal Soap.....	p.104
Vendex 50WP.....	p.122
Vydate L.....	p.125

Nematodes

Vydate L.....	p.126
---------------	-------

Plant Bugs

Sevin 4F.....	p.110
---------------	-------

Spider Mites

Dibrom 8 Emulsive.....	p.31
Malathion 5 EC.....	p.77
Malathion 8	
AQUAMUL.....	p.83

Stink Bugs

Sevin 4F.....	p.110
Sevin SL.....	p.117

Thrips

Sevin 4F.....	p.110
Sevin SL.....	p.117

Whiteflies

Alias 4F.....	p.19
Safer Insecticidal Soap.....	p.104

Diseases

Alternaria Blight

Kocide 3000.....p.63

Anthracnose

Equis 720.....p.54

Kocide 3000.....p.63

Botrytis Blight (Gray Mold)

Equis 720.....p.54

Cercospora Leaf Spot

Equis 720.....p.54

Powdery Mildew

Equis 720.....p.54

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Garlic

Insects, Mites and Nematodes

Aphids

Malathion 5 EC.....p.78
 Malathion 8

AQUAMUL.....p.83

Armyworms

Dipel DF.....p.38
 Lannate LV.....p.72
 Perm-Up 3.2 EC.....p.98

Caterpillars

Dipel DF.....p.38

Cutworms

Dipel DF.....p.38
 Perm-Up 3.2 EC.....p.98

Diamondback Moths

Dipel DF.....p.38

Leafminers

Perm-Up 3.2 EC.....p.98

Leafrollers

Dipel DF.....p.38

Loopers

Dipel DF.....p.38

Stink Bugs

Perm-Up 3.2 EC.....p.98

Thrips

Malathion 5 EC.....p.78
 Malathion 8

AQUAMUL.....p.83

Perm-Up 3.2 EC.....p.98

Safer Insecticidal Soap.....p.105

Diseases

Bacterial Blight

Kocide 3000.....p.64

Botrytis Leaf Blight/Blast (Gray Mold)

Bravo 720.....p.25
 Equis 720.....p.55
 Manzate Pro Stick.....p.91

Botrytis Neck Rot

Equis 720.....p.55

Downy Mildew

Equis 720.....p.55
 Kocide 3000.....p.64
 Manzate Pro Stick.....p.91

Neck Rot

Manzate Pro Stick.....p.91

Purple Blotch

Equis 720.....p.55
 Kocide 3000.....p.64
 Manzate Pro Stick.....p.91

Other

Slugs and Snails

CROP AND PEST USE INDEX

Sluggo Snail & Slug Bait.....p.120

Gourds

Insects, Mites and Nematodes

Aphids

Lannate LV.....p.74

Armyworms

Dipel DF.....p.36

Lannate LV.....p.74

Cucumber Beetles

Lannate LV.....p.74

Cutworms

Lannate LV.....p.74

Caterpillars

Lannate LV.....p.74

Flea Beetles

Lannate LV.....p.74

Loopers

Dipel DF.....p.36

Lannate LV.....p.74

Melonworms

Dipel DF.....p.36

Lannate LV.....p.74

Diseases

Anthracnose

Equis 720.....p.52

Downy Mildew

Equis 720.....p.52

Gummy Stem Blight/Vine Decline

Equis 720.....p.53

Powdery Mildew (Sphaerotheca only)

Equis 720.....p.53

Target Spot

Equis 720.....p.52

Scab

Equis 720.....p.53

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Grapefruit

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Safer Insecticidal Soap.....	p.103
Vydate L.....	p.125

Beetles

Sevin 4F.....	p.108
Vydate L.....	p.125

Black Flies

Alias 4F.....	p.18
---------------	------

Cutworms

Dipel DF.....	p.37
Sevin 4F.....	p.107
Sevin SL.....	p.116

Flea Beetles

Alias 4F.....	p.18
---------------	------

Leafhoppers

Alias 4F.....	p.18
Vydate L.....	p.125

Leafminers

Alias 4F.....	p.18
Vydate L.....	p.125

Leafrollers

Dibrom 8 Emulsive.....	p.31
Dipel DF.....	p.37
Sevin 4F.....	p.107
Sevin SL.....	p.116

Mealybugs

Alias 4F.....	p.18
Vydate L.....	p.125

Mites

Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Safer Insecticidal Soap.....	p.103
Sevin 4F.....	p.107
Sevin SL.....	p.116
Vendex 50WP.....	p.122
Vydate L.....	p.124/125

Nematodes

Vydate L.....	p.125
---------------	-------

Plant Bugs

Sevin 4F.....	p.107
Sevin SL.....	p.116

Scale Crawlers

Dibrom 8 Emulsive.....	p.31
------------------------	------

Scales

Alias 4F.....	p.18
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Malathion 8	
AQUAMUL.....	p.82
Safer Insecticidal Soap.....	p.103
Sevin 4F.....	p.107
Sevin SL.....	p.116
Vydate L.....	p.125

Sharpshooters

Alias 4F.....	p.18
---------------	------

Thrips

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.34
Malathion 5 EC.....	p.77
Vydate L.....	p.124/125

Weevils

Sevin 4F.....p.108

Sevin SL.....p.117

Whiteflies

Alias 4F.....p.18

Dimethoate 4 EC.....p.34

Diseases

Alternaria

Kocide 3000.....p.60

Black spot

Kocide 3000.....p.62

Brown Rot

Kocide 3000.....p.61

Brown spot

Kocide 3000.....p.60

Citrus Canker

Kocide 3000.....p.61

Melanose

Kocide 3000.....p.60

Phytophthora

Kocide 3000.....p.61

Phytophthora Foot Rot

Kocide 3000.....p.61

Scab

Kocide 3000.....p.60

Septoria Spot

Kocide 3000.....p.61

Other

Slugs and Snails

Deadline GT.....p.48

Sluggo Snail & Slug Bait.....p.120

Green Onion

Insects, Mites and Nematodes

Armyworms

Dipel DF.....p.38

Lannate LV.....p.73

Caterpillars

Dipel DF.....p.38

Cutworms

Dipel DF.....p.38

Lannate LV.....p.73

Diamondback Moths

Dipel DF.....p.38

Leafrollers

Dipel DF.....p.38

Loopers

Dipel DF.....p.38

Thrips

Lannate LV.....p.73

Diseases

Botrytis Blight (Gray Mold)

Bravo 720.....p.25

Guava

Insects, Mites and Nematodes

Ants

Amdro Fire Ant Bait.....p.22

Aphids

Alias 4F.....p.21

Leafhoppers

Alias 4F.....p.21

Mealybugs

Alias 4F.....p.21

Scales

Alias 4F.....p.21

Thrips

Alias 4F.....p.21

Whiteflies

Alias 4F.....p.21

Diseases

Anthracnose

Kocide 3000.....p.63

Red Algae

Kocide 3000.....p.63

Honeydew

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.73
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97
Safer Insecticidal Soap.....	p.104
Vydate L.....	p.123

Armyworms

Dipel DF.....	p.36
Lannate LV.....	p.73

Caterpillars

Lannate LV.....	p.73
-----------------	------

Cucumber Beetles

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.73
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97

Cutworms

D.Z.N Diazinon.....	p.41
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.97

Flea Beetles

Lannate LV.....	p.73
-----------------	------

Leafhoppers

Alias 4F.....	p.18
Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.42
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97

Leafminers

Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.42
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97
Vydate L.....	p.123

Loopers

Dipel DF.....	p.36
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.97

Melonworms

Dipel DF.....	p.36
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.97

Mites

Safer Insecticidal Soap.....	p.104
------------------------------	-------

Nematodes

Vydate L.....	p.123
---------------	-------

Plant Bugs

Perm-Up 3.2 EC.....	p.97
Safer Insecticidal Soap.....	p.104

Spider Mites

D.Z.N Diazinon.....	p.42
Malathion 5 EC.....	p.78
Manzate Pro Stick.....	p.91

Squash Bugs

Perm-Up 3.2 EC.....	p.97
---------------------	------

Thrips

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.42

Whiteflies

Alias 4F.....	p.18
Safer Insecticidal Soap.....	p.104

Diseases

Alternaria Leaf Blight

Bravo 720.....p.24
Equis 720.....p.52

Alternaria Leaf Spot

Equis 720.....p.52
Kocide 3000.....p.63
Manzate Pro Stick.....p.91

Angular Leaf Spot

Kocide 3000.....p.63

Anthracnose

Bravo 720.....p.24
Equis 720.....p.52
Kocide 3000.....p.63
Manzate Pro Stick.....p.91

Cercospora Leaf Spot

Bravo 720.....p.24
Equis 720.....p.52
Manzate Pro Stick.....p.91

Downy Mildew

Equis 720.....p.52
Kocide 3000.....p.63
Manzate Pro Stick.....p.91

Gummy Stem Blight

Bravo 720.....p.24
Equis 720.....p.52
Kocide 3000.....p.63

Powdery Mildew

Bravo 720.....p.24
Equis 720.....p.52

Scab

Bravo 720.....p.24
Equis 720.....p.52

Target Spot

Bravo 720.....p.24
Equis 720.....p.52

Watermelon Bacterial Fruit Blotch

Kocide 3000.....p.63

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Horseradish

Insects, Mites and Nematodes

Aphids

Lannate LV.....p.72

Malathion 5 EC.....p.80

Malathion 8

AQUAMUL.....p.85

Armyworms

Sevin 4F.....p.114

Beetles

Sevin 4F.....p.114

Caterpillars

Sevin 4F.....p.114

Cutworms

Sevin 4F.....p.114

Flea Beetles

Sevin 4F.....p.113

Leafhoppers

Sevin 4F.....p.113

Plant Bugs

Sevin 4F.....p.114

Stink Bugs

Sevin 4F.....p.114

Thrips

Lannate LV.....p.72

Weevils

Perm-Up 3.2 EC.....p.98

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Iba

Consult your agricultural extension agent for information on pesticide use for this crop.

Jackfruit

Consult your agricultural extension agent for information on pesticide use for this crop.

Kondot

Insects, Mites and Nematodes

Aphids

Lannate LV.....p.74

Armyworms

Dipel DF.....p.36

Lannate LV.....p.74

Caterpillars

Lannate LV.....p.74

Cucumber Beetles

Lannate LV.....p.74

Cutworms

Lannate LV.....p.74

Flea Beetles

Lannate LV.....p.74

Loopers

Dipel DF.....p.36

Lannate LV.....p.74

Melonworms

Dipel DF.....p.36

Lannate LV.....p.74

Lemon

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Safer Insecticidal Soap.....	p.103
Vydate L.....	p.125

Beetles

Sevin 4F.....	p.108
Vydate L.....	p.125
Alias 4F.....	p.18

Cutworms

Dipel DF.....	p.37
Sevin 4F.....	p.107
Sevin SL.....	p.116

Flea Beetles

Alias 4F.....	p.18
---------------	------

Leafhoppers

Alias 4F.....	p.18
Vydate L.....	p.125

Leafminers

Alias 4F.....	p.18
Vydate L.....	p.125

Leafrollers

Dibrom 8 Emulsive.....	p.31
Dipel DF.....	p.37
Sevin 4F.....	p.107
Sevin SL.....	p.116

Mealybugs

Alias 4F.....	p.18
Vydate L.....	p.125

Mites

Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Safer Insecticidal Soap.....	p.103
Sevin 4F.....	p.107
Sevin SL.....	p.116
Vendex 50WP.....	p.122
Vydate L.....	p.124/125

Nematodes

Vydate L.....	p.125
---------------	-------

Plant Bugs

Sevin 4F.....	p.107
Sevin SL.....	p.116

Scale Crawlers

Dibrom 8 Emulsive.....	p.31
------------------------	------

Scales

Alias 4F.....	p.18
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Malathion 8	
AQUAMUL.....	p.82
Safer Insecticidal Soap.....	p.103
Sevin 4F.....	p.107
Sevin SL.....	p.116
Vydate L.....	p.125

Sharpshooters

Alias 4F.....	p.18
---------------	------

Thrips

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Vydate L.....	p.124/125

Weevils

Sevin 4F.....p.108

Sevin SL.....p.117

Whiteflies

Alias 4F.....p.18

Dimethoate 4 EC.....p.33

Vydate L.....p.125

Diseases

Alternaria

Kocide 3000.....p.60

Black Spot

Kocide 3000.....p.62

Brown Rot

Kocide 3000.....p.61

Brown Spot

Kocide 3000.....p.60

Citrus Canker

Kocide 3000.....p.61

Melanose

Kocide 3000.....p.60

Phytophthora

Kocide 3000.....p.61

Phytophthora Foot Rot

Kocide 3000.....p.61

Scab

Kocide 3000.....p.60

Septoria Spot

Kocide 3000.....p.61

Other

Slugs and Snails

Deadline GT.....p.48

Sluggo Snail & Slug Bait.....p.119

Lettuce, Head

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.19
Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.43
Lannate LV.....	p.72
Perm-Up 3.2 EC.....	p.99
Safer Insecticidal Soap.....	p.105

Armyworms

Dipel DF.....	p.39
Lannate LV.....	p.72
Perm-Up 3.2 EC.....	p.99
Sevin 4F.....	p.111
Sevin SL.....	p.118

Caterpillars

Dipel DF.....	p.39
Lannate LV.....	p.72
Sevin 4F.....	p.111

Cutworms

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.43
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.99

Diamondback Moths

Dipel DF.....	p.39
---------------	------

Flea Beetles

Alias 4F.....	p.19
Sevin 4F.....	p.111
Sevin SL.....	p.118

Leafhoppers

Alias 4F.....	p.19
Dimethoate 4 EC.....	p.34
Lannate LV.....	p.72
Perm-Up 3.2 EC.....	p.99
Sevin 4F.....	p.111
Sevin SL.....	p.118

Leafminers

Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.43
Perm-Up 3.2 EC.....	p.99

Leafrollers

Dipel DF.....	p.39
---------------	------

Loopers

Dipel DF.....	p.39
Lannate LV.....	p.72
Perm-Up 3.2 EC.....	p.99

Mites

Safer Insecticidal Soap.....	p.105
------------------------------	-------

Plant Bugs

Sevin 4F.....	p.111
Sevin SL.....	p.118

Stink Bugs

Sevin 4F.....	p.111
Sevin SL.....	p.118

Thrips

Lannate LV.....	p.72
-----------------	------

Whiteflies

Alias 4F.....	p.19
Safer Insecticidal Soap.....	p.105

Diseases

Anthracnose

Manzate Pro Stick.....p.90

Downy Mildew

Kocide 3000.....p.64

Manzate Pro Stick.....p.90

Other

Slugs and Snails

Deadline GT.....p.48

Sluggo Snail & Slug Bait.....p.120

Lettuce, Leaf

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.19
Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.43
Lannate LV.....	p.72
Malathion 5 EC.....	p.78
Malathion 8 AQUAMUL.....	

p.83

Perm-Up 3.2 EC.....	p.99
Safer Insecticidal Soap.....	p.105

Armyworms

Dipel DF.....	p.39
Lannate LV.....	p.72
Perm-Up 3.2 EC.....	p.99
Sevin 4F.....	p.111
Sevin SL.....	p.118

Caterpillars

Dipel DF.....	p.39
Lannate LV.....	p.72
Sevin 4F.....	p.111

Cutworms

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.43
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.99

Diamondback Moths

Dipel DF.....	p.39
---------------	------

Flea Beetles

Alias 4F.....	p.19
Sevin 4F.....	p.111
Sevin SL.....	p.118

Leafhoppers

Alias 4F.....	p.19
Dimethoate 4 EC.....	p.35
Lannate LV.....	p.72
Malathion 5 EC.....	p.78
Malathion 8 AQUAMUL.....	

p.83

Perm-Up 3.2 EC.....	p.99
Sevin 4F.....	p.111
Sevin SL.....	p.118

Leafminers

Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.43
Perm-Up 3.2 EC.....	p.99

Leafrollers

Dipel DF.....	p.39
---------------	------

Loopers

Dipel DF.....	p.39
Lannate LV.....	p.72
Malathion 5 EC.....	p.78
Malathion 8 AQUAMUL.....	

p.84

Perm-Up 3.2 EC.....	p.99
---------------------	------

Mites

Safer Insecticidal Soap.....	p.105
------------------------------	-------

Plant Bugs

Sevin 4F.....	p.111
Sevin SL.....	p.118

Scales

Malathion 5 EC.....	p.78
---------------------	------

Stink Bugs

Sevin 4F.....	p.111
Sevin SL.....	p.118

CROP AND PEST USE INDEX

Spider Mites

Malathion 5 EC.....p.78
Malathion 8 AQUAMUL.....

p.84

Thrips

Lannate LV.....p.72
Malathion 5 EC.....p.78

Whiteflies

Alias 4F.....p.19
Safer Insecticidal Soap.....p.105

Diseases

Anthracnose

Manzate Pro Stick.....p.90

Downy Mildew

Kocide 3000.....p.64
Manzate Pro Stick.....p.90

Other

Slugs and Snails

Deadline GT.....p.48
Sluggo Snail & Slug Bait.....p.120

Lime

Insects, Mites and Nematodes

Aphids

Alias 4F.....p.18
Vydate L.....p.125

Beetles

Sevin 4F.....p.108
Vydate L.....p.125

Cutworms

Alias 4F.....p.18
Dipel DF.....p.37
Sevin 4F.....p.107
Sevin SL.....p.116

Flea Beetles

Alias 4F.....p.18

Leafhoppers

Alias 4F.....p.18
Vydate L.....p.125

Leafminers

Alias 4F.....p.18

Leafrollers

Dipel DF.....p.37
Sevin 4F.....p.107
Sevin SL.....p.116

Mealybugs

Alias 4F.....p.18
Vydate L.....p.125

Mites

Malathion 5 EC.....p.77
Sevin 4F.....p.107
Sevin SL.....p.116
Vendex 50WP.....p.122
Vydate L.....p.124/125

Nematodes

Vydate L.....p.125

Plant Bugs

Sevin 4F.....p.107
Sevin SL.....p.116

Scales

Alias 4F.....p.18
Malathion 5 EC.....p.77
Malathion 8
AQUAMUL.....p.82
Sevin 4F.....p.107
Sevin SL.....p.116
Vydate L.....p.125

Sharpshooters

Alias 4F.....p.18

Thrips

Alias 4F.....p.18
Malathion 5 EC.....p.77
Vydate L.....p.124/125

Weevils

Sevin 4F.....p.108
Sevin SL.....p.117

Whiteflies

Alias 4F.....p.18
Vydate L.....p.125

Diseases

Alternaria

Kocide 3000.....p.60

Black Spot

Kocide 3000.....p.62

Brown Rot

Kocide 3000.....p.61

Brown Spot

Kocide 3000.....p.60

Citrus Canker

Kocide 3000.....p.61

Melanose

Kocide 3000.....p.60

Phytophthora

Kocide 3000.....p.61

Phytophthora Foot Rot

Kocide 3000.....p.61

Scab

Kocide 3000.....p.60

Septoria Spot

Kocide 3000.....p.61

Other

Slugs and Snails

Deadline GT.....p.48

Sluggo Snail & Slug Bait.....p.119

Mango

Insects, Mites and Nematodes

Ants

Amdro Fire Ant Bait.....p.22

Aphids

Alias 4F.....p.21

Leafhoppers

Alias 4F.....p.21

Mealybugs

Alias 4F.....p.21

Scales

Alias 4F.....p.21

Thrips

Alias 4F.....p.21

Whiteflies

Alias 4F.....p.21

Diseases

Anthracnose

Equis 720.....p.54

Kocide 3000.....p.64

Manzate Pro Stick.....p.93

Black Spot (*Cercospora*)

Manzate Pro Stick.....p.93

Phytophthora Fruit Rot

Manzate Pro Stick.....p.93

Muskmelon

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.73
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97
Safer Insecticidal Soap.....	p.104

Armyworms

Dibrom 8 Emulsive.....	p.30
Dipel DF.....	p.36
Lannate LV.....	p.73

Caterpillars

Lannate LV.....	p.73
-----------------	------

Cucumber Beetles

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.73
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97

Cutworms

D.Z.N Diazinon.....	p.41
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.97

Flea Beetles

Lannate LV.....	p.73
-----------------	------

Leafhoppers

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.42
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97

Leafminers

Dibrom 8 Emulsive.....	p.30
Dimethoate 4 EC.....	p.34
D.Z.N Diazinon.....	p.42
Malathion 5 EC.....	p.78
Perm-Up 3.2 EC.....	p.97

Loopers

Dibrom 8 Emulsive.....	p.30
Dipel DF.....	p.36
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.97

Melonworms

Dipel DF.....	p.36
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.73
Perm-Up 3.2 EC.....	p.97

Mites

Safer Insecticidal Soap.....	p.104
------------------------------	-------

Plant Bugs

Perm-Up 3.2 EC.....	p.97
Safer Insecticidal Soap.....	p.104

Squash Bugs

Perm-Up 3.2 EC.....	p.97
---------------------	------

Spider Mites

D.Z.N Diazinon.....	p.42
Malathion 5 EC.....	p.78

Thrips

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.42

Whiteflies

Alias 4F.....	p.18
Safer Insecticidal Soap.....	p.104

Diseases

Alternaria Leaf Blight

Bravo 720.....p.25

Alternaria Leaf Spot

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Angular Leaf Spot

Kocide 3000.....p.63

Anthracnose

Bravo 720.....p.24

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Cercospora Leaf Spot

Bravo 720.....p.25

Manzate Pro Stick.....p.90

Downy Mildew

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Gummy Stem Blight

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Powdery Mildew

Bravo 720.....p.25

Kocide 3000.....p.63

Scab

Manzate Pro Stick.....p.90

Target Spot

Bravo 720.....p.24

Equis 720.....p.52

Watermelon Bacterial Fruit Blotch

Kocide 3000.....p.63

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Mustard Greens

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.17
Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.44

Armyworms

Dipel DF.....	p.39
Lannate LV.....	p.72
Sevin 4F.....	p.106
Sevin SL.....	p.116

Caterpillars

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.44
Lannate LV.....	p.72
Sevin 4F.....	p.106

Cutworms

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.43

Diamondback Moths

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.44
Lannate LV.....	p.72
Sevin 4F.....	p.106
Sevin SL.....	p.116

Flea Beetles

Alias 4F.....	p.17
Sevin 4F.....	p.106
Sevin SL.....	p.115

Leafhoppers

Alias 4F.....	p.17
Dimethoate 4 EC.....	p.35
Sevin 4F.....	p.106
Sevin SL.....	p.115

Leafminers

Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.44

Leafrollers

Dipel DF.....	p.39
---------------	------

Loopers

Dipel DF.....	p.39
Lannate LV.....	p.72

Plant Bugs

Sevin 4F.....	p.106
Sevin SL.....	p.116

Stink Bugs

Sevin 4F.....	p.106
Sevin SL.....	p.116

Whiteflies

Alias 4F.....	p.17
---------------	------

Diseases

Alternaria Leaf Spot

Equis 720.....p.51

Black Leaf Spot (Alternaria)

Kocide 3000.....p.62

Black Rot (Xanthomonas)

Kocide 3000.....p.62

Downy Mildew

Equis 720.....p.51

Kocide 3000.....p.62

Other

Slugs and Snails

Deadline GT.....p.48

Sluggo Snail & Slug Bait.....p.120

Okra

Insects, Mites and Nematodes

Aphids

Alias 4F.....p.19
 Malathion 5 EC.....p.78
 Malathion 8

AQUAMUL.....p.84

Beetles

Malathion 8

AQUAMUL.....p.84

Colorado Potato Beetle

Alias 4F.....p.19

Cucumber Beetles

Malathion 5 EC.....p.78

Leafhoppers

Alias 4F.....p.19

Malathion 5 EC.....p.78

Leafminers

Malathion 5 EC.....p.78

Spider Mites

Malathion 5 EC.....p.78

Stink Bugs

Sevin 4F.....p.113

Whiteflies

Alias 4F.....p.19

Diseases

Anthracnose

Equis 720.....p.54
 Kocide 3000.....p.64

Bacterial Leaf Spot

Kocide 3000.....p.64

Botrytis Blight (Gray Mold)

Equis 720.....p.54

Cercospora Leaf Spot

Equis 720.....p.54

Leaf Spot

Kocide 3000.....p.64

Pod Spot

Kocide 3000.....p.64

Powdery Mildew

Equis 720.....p.54
 Kocide 3000.....p.64

Other

Slugs and Snails

CROP AND PEST USE INDEX

Sluggo Snail & Slug Bait.....p.120

Orange

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Safer Insecticidal Soap.....	p.103
Vydate L.....	p.125

Beetles

Sevin 4F.....	p.108
Vydate L.....	p.125

Black Flies

Alias 4F.....	p.18
---------------	------

Cutworms

Dipel DF.....	p.37
Sevin 4F.....	p.107
Sevin SL.....	p.116

Flea Beetles

Alias 4F.....	p.18
---------------	------

Leafhoppers

Alias 4F.....	p.18
Vydate L.....	p.125

Leafminers

Alias 4F.....	p.18
Vydate L.....	p.125

Leafrollers

Dibrom 8 Emulsive.....	p.31
Dipel DF.....	p.37
Sevin 4F.....	p.107
Sevin SL.....	p.116

Mealybugs

Alias 4F.....	p.18
Vydate L.....	p.125

Mites

Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Safer Insecticidal Soap.....	p.103
Sevin 4F.....	p.107
Sevin SL.....	p.116
Vendex 50WP.....	p.122
Vydate L.....	p.124/125

Nematodes

Vydate L.....	p.125
---------------	-------

Plant Bugs

Sevin 4F.....	p.107
Sevin SL.....	p.116

Scale Crawlers

Dibrom 8 Emulsive.....	p.31
------------------------	------

Scales

Alias 4F.....	p.18
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Malathion 8	
AQUAMUL.....	p.82
Safer Insecticidal Soap.....	p.103
Sevin 4F.....	p.107
Sevin SL.....	p.116
Vydate L.....	p.125

Sharpshooters

Alias 4F.....	p.18
---------------	------

Thrips

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Vydate L.....	p.124/125

Weevils

Sevin 4F.....p.108
Sevin SL.....p.117

Whiteflies

Alias 4F.....p.18
Dimethoate 4 EC.....p.33
Vydate L.....p.125

Diseases

Alternaria

Kocide 3000.....p.60

Black Spot

Kocide 3000.....p.62

Brown Rot

Kocide 3000.....p.61

Brown Spot

Kocide 3000.....p.60

Citrus Canker

Kocide 3000.....p.61

Melanose

Kocide 3000.....p.60

Phytophthora

Kocide 3000.....p.61

Phytophthora Foot Rot

Kocide 3000.....p.61

Scab

Kocide 3000.....p.60

Septoria Spot

Kocide 3000.....p.61

Other

Slugs and Snails

Deadline GT.....p.48
Sluggo Snail & Slug Bait.....p.119

Papaya

Insects, Mites and Nematodes

Ants

Amdro Fire Ant Bait.....p.22

Aphids

Alias 4F.....p.21

Malathion 5 EC.....p.79

Malathion 8

AQUAMUL.....p.84

Leafhoppers

Alias 4F.....p.21

Mealybugs

Alias 4F.....p.21

Malathion 5 EC.....p.79

Malathion 8

AQUAMUL.....p.84

Mites

Vendex 50WP.....p.122

Scales

Alias 4F.....p.21

Thrips

Alias 4F.....p.21

Whiteflies

Alias 4F.....p.21

Diseases

Alternaria Fruit Spot

Equis 720.....p.55

Anthracnose

Bravo 720.....p.26

Equis 720.....p.55

Kocide 3000.....p.63

Manzate Pro Stick.....p.92

Black Spot (Cercospora)

Manzate Pro Stick.....p.92

Phytophthora Fruit Rot

Manzate Pro Stick.....p.92

Red Algae

Kocide 3000.....p.63

Stem End Rot

Bravo 720.....p.26

Equis 720.....p.55

Parsley

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.19
Malathion 5 EC.....	p.79
Malathion 8	
AQUAMUL.....	p.84
Perm-Up 3.2 EC.....	p.99

Armyworms

Dipel DF.....	p.39
Lannate LV.....	p.72
Perm-Up 3.2 EC.....	p.99
Sevin 4F.....	p.111
Sevin SL.....	p.118

Caterpillars

Dipel DF.....	p.39
Lannate LV.....	p.72
Sevin 4F.....	p.111

Colorado Potato Beetle

Alias 4F.....	p.19
---------------	------

Cutworms

Dipel DF.....	p.39
D.Z.N Diazinon.....	p.44
Perm-Up 3.2 EC.....	p.99

Diamondback Moths

Dipel DF.....	p.39
Lannate LV.....	p.72

Flea Beetles

Alias 4F.....	p.20
Sevin 4F.....	p.111

Sevin SL.....	p.118
---------------	-------

Leafhoppers

Alias 4F.....	p.19/20
Perm-Up 3.2 EC.....	p.99
Sevin 4F.....	p.111
Sevin SL.....	p.118

Leafminers

Perm-Up 3.2 EC.....	p.99
---------------------	------

Leafrollers

Dipel DF.....	p.39
---------------	------

Loopers

Dipel DF.....	p.39
Lannate LV.....	p.72
Perm-Up 3.2 EC.....	p.99

Plant Bugs

Sevin 4F.....	p.111
Sevin SL.....	p.118

Stink Bugs

Sevin 4F.....	p.111
Sevin SL.....	p.118

Whiteflies

Alias 4F.....	p.19/20
---------------	---------

Diseases

Bacterial Blight (*Pseudomonas*)

CROP AND PEST USE INDEX

Kocide 3000.....p.65

Peanut

Insects, Mites and Nematodes

Aphids

Alias 4F.....p.20

Armyworms

Lannate LV.....p.74

Caterpillars

Dipel DF.....p.39

Lannate LV.....p.74

Cutworms

Lannate LV.....p.74

Leafhoppers

Alias 4F.....p.20

Lannate LV.....p.74

Loopers

Dipel DF.....p.39

Lannate LV.....p.74

Thrips

Alias 4F.....p.20

Lannate LV.....p.74

Whiteflies

Alias 4F.....p.20

Diseases

Ascochyta Web Blotch

Manzate Pro Stick.....p.92

Early Leaf Spot (Cercospora)

Bravo 720.....p.26

Equis 720.....p.56

Kocide 3000.....p.65

Manzate Pro Stick.....p.92

Late Leaf Spot (Cercosporidium)

Bravo 720.....p.26

Equis 720.....p.56

Pepper Spot

Equis 720.....p.56

Rust

Bravo 720.....p.26

Equis 720.....p.56

Manzate Pro Stick.....p.92

Web Blotch

Equis 720.....p.56

Pepino

Insects, Mites and Nematodes

Aphids

- Alias 4F.....p.19
- Dimethoate 4 EC.....p.34
- Lannate LV.....p.73
- Safer Insecticidal Soap.....p.104

Armyworms

- Dipel DF.....p.36
- Lannate LV.....p.73
- Sevin SL.....p.117

Caterpillars

- Lannate LV.....p.73

Colorado Potato Beetle

- Alias 4F.....p.19

Cucumber Beetles

- Lannate LV.....p.73

Cutworms

- Lannate LV.....p.73
- Sevin SL.....p.118

Flea Beetles

- Lannate LV.....p.73
- Sevin SL.....p.117

Leafhoppers

- Alias 4F.....p.19
- Dimethoate 4 EC.....p.34
- Sevin SL.....p.117

Leafminers

- Dimethoate 4 EC.....p.34

Loopers

- Dipel DF.....p.36
- Lannate LV.....p.73

Melonworms

- Dipel DF.....p.36
- Lannate LV.....p.73

Mites

- Safer Insecticidal Soap.....p.104

Plant Bugs

- Safer Insecticidal Soap.....p.104

Stink Bugs

- Sevin SL.....p.117

Thrips

- Sevin SL.....p.117

Whiteflies

- Alias 4F.....p.19
- Safer Insecticidal Soap.....p.104

Diseases

Anthracnose

- Equis 720.....p.54

Botrytis Blight (Gray Mold)

- Equis 720.....p.54

Cercospora Leaf Spot

- Equis 720.....p.54

Powdery Mildew

- Equis 720.....p.54

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Pepper, Bell

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.19
Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.44
Lannate LV.....	p.74
Malathion 5 EC.....	p.79
Malathion 8	
AQUAMUL.....	p.84
Safer Insecticidal Soap.....	p.104
Vydate L.....	p.126

Armyworms

Dipel DF.....	p.38
Lannate LV.....	p.74
Sevin 4F.....	p.110
Sevin SL.....	p.117

Beetles

Dibrom 8 Emulsive.....	p.31
Sevin 4F.....	p.110

Caterpillars

Dipel DF.....	p.38
Lannate LV.....	p.74
Malathion 5 EC.....	p.79
Sevin 4F.....	p.110

Colorado Potato Beetle

Alias 4F.....	p.19
---------------	------

Cutworms

Dipel DF.....	p.38
D.Z.N Diazinon.....	p.44
Lannate LV.....	p.74
Perm-Up 3.2 EC.....	p.100
Sevin 4F.....	p.111
Sevin SL.....	p.118

Flea Beetles

Dibrom 8 Emulsive.....	p.31
Perm-Up 3.2 EC.....	p.100
Sevin 4F.....	p.110
Sevin SL.....	p.117

Leafhoppers

Alias 4F.....	p.19
Dimethoate 4 EC.....	p.35
Sevin 4F.....	p.110
Sevin SL.....	p.117

Leafminers

Dibrom 8 Emulsive.....	p.31
D.Z.N Diazinon.....	p.44
Perm-Up 3.2 EC.....	p.100
Vydate L.....	p.126

Loopers

Dipel DF.....	p.38
Lannate LV.....	p.74
Perm-Up 3.2 EC.....	p.100

Mites

Safer Insecticidal Soap.....	p.104
------------------------------	-------

Nematodes

Vydate L.....	p.126
---------------	-------

Plant Bugs

Sevin 4F.....	p.110
---------------	-------

Spider Mites

Dibrom 8 Emulsive.....	p.31
------------------------	------

Stink Bugs

Sevin 4F.....	p.110
Sevin SL.....	p.117

Thrips

Sevin 4F.....	p.110
Sevin SL.....	p.117

CROP AND PEST USE INDEX

Weevils

Perm-Up 3.2 EC.....p.100
Vydate L.....p.126

Safer Insecticidal Soap.....p.104

Whiteflies

Alias 4F.....p.19

Diseases

Anthracnose

Equis 720.....p.54
Kocide 3000.....p.65

Cercospora Leaf Spot

Equis 720.....p.54
Kocide 3000.....p.65

Bacterial Spot

Kocide 3000.....p.65

Powdery Mildew

Equis 720.....p.54

Botrytis Blight (Gray Mold)

Equis 720.....p.54

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Pepper, Hot

Insects, Mites and Nematodes

Aphids

Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.44
Lannate LV.....	p.74
Malathion 8 AQUAMUL.....	p.
Malathion 5 EC.....	p.79
Safer Insecticidal Soap.....	p.104
Vydate L.....	p.126

Armyworms

Dipel DF.....	p.38
Lannate LV.....	p.74
Sevin 4F.....	p.110

Beetles

Sevin 4F.....	p.110
---------------	-------

Caterpillars

Dipel DF.....	p.38
Lannate LV.....	p.74
Malathion 5 EC.....	p.79
Sevin 4F.....	p.110

Cutworms

Dipel DF.....	p.38
D.Z.N Diazinon.....	p.44
Lannate LV.....	p.74
Sevin 4F.....	p.111

Flea Beetles

Sevin 4F.....	p.110
---------------	-------

Leafhoppers

Sevin 4F.....	p.110
---------------	-------

Leafminers

Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.44
Vydate L.....	p.126

Leafrollers

Dibrom 8 Emulsive.....	p.31
------------------------	------

Loopers

Dipel DF.....	p.38
Lannate LV.....	p.74

Mites

Dibrom 8 Emulsive.....	p.31
Safer Insecticidal Soap.....	p.104

Nematodes

Vydate L.....	p.126
---------------	-------

Plant Bugs

Sevin 4F.....	p.110
---------------	-------

Scale Crawlers

Dibrom 8 Emulsive.....	p.31
------------------------	------

Stink Bugs

Sevin 4F.....	p.110
---------------	-------

Thrips

Dibrom 8 Emulsive.....	p.31
Sevin 4F.....	p.110

Weevils

Vydate L.....	p.126
---------------	-------

Whiteflies

Safer Insecticidal Soap.....	p.104
------------------------------	-------

Diseases

Anthracnose

Kocide 3000.....p.65

Cercospora Leaf Spot

Kocide 3000.....p.65

Bacterial Spot

Kocide 3000.....p.65

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Pineapple

Insects, Mites and Nematodes

Mealybugs

Malathion 5 EC.....p.79

Malathion 8

AQUAMUL.....p.85

Nematodes

Vydate L.....p.126

Pomegranate

Consult your agricultural extension for information on pesticide use for this crop.

Pumpkin

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
Malathion 8	
AQUAMUL.....	p.85
Perm-Up 3.2 EC.....	p. 97
Vydate L.....	p.123

Armyworms

Dipel DF.....	p.36
---------------	------

Caterpillars

Malathion 8	
AQUAMUL.....	p.85
Sevin 4F.....	p.109

Cucumber Beetles

Alias 4F.....	p.18
Malathion 8	
AQUAMUL.....	p.85
Perm-Up 3.2 EC.....	p. 97
Sevin 4F.....	p.110

Cutworms

Perm-Up 3.2 EC.....	p. 97
---------------------	-------

Flea Beetles

Sevin 4F.....	p.110
---------------	-------

Leafhoppers

Alias 4F.....	p.18
Malathion 8	
AQUAMUL.....	p.85
Perm-Up 3.2 EC.....	p. 97
Sevin 4F.....	p.110

Leafminers

Perm-Up 3.2 EC.....	p. 97
---------------------	-------

Vydate L.....	p.123
---------------	-------

Loopers

Dipel DF.....	p.36
Perm-Up 3.2 EC.....	p. 97

Melonworms

Dipel DF.....	p.36
Perm-Up 3.2 EC.....	p. 97
Sevin 4F.....	p.109

Nematodes

Vydate L.....	p.123
---------------	-------

Plant Bugs

Perm-Up 3.2 EC.....	p. 97
---------------------	-------

Squash Bugs

Perm-Up 3.2 EC.....	p. 97
Sevin 4F.....	p.110

Spider Mites

Malathion 8	
AQUAMUL.....	p.85

Thrips

Alias 4F.....	p.18
Malathion 8	
AQUAMUL.....	p.85

Whiteflies

Alias 4F.....	p.18
---------------	------

Diseases

Alternaria Leaf Blight

Bravo 720.....p.25

Alternaria Leaf Spot

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Angular Leaf Spot

Kocide 3000.....p.63

Anthracnose

Bravo 720.....p.24

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Cercospora Leaf Spot

Bravo 720.....p.25

Manzate Pro Stick.....p.90

Downy Mildew

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Gummy Stem Blight

Bravo 720.....p.25

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Powdery Mildew

Bravo 720.....p.25

Kocide 3000.....p.63

Scab

Bravo 720.....p.25

Manzate Pro Stick.....p.90

Target Spot

Bravo 720.....p.24

Equis 720.....p.52

Watermelon Bacterial Fruit Blotch

Kocide 3000.....p.63

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Radish

Insects, Mites and Nematodes

Aphids

Alias 4F.....p.20
 D.Z.N Diazinon.....p.44
 Malathion 5 EC.....p.80
 Malathion 8

AQUAMUL.....p.85

Safer Insecticidal Soap.....p.103

Armyworms

Sevin 4F.....p.114

Beetles

Sevin 4F.....p.114

Caterpillars

Sevin 4F.....p.114

Cutworms

D.Z.N Diazinon.....p.44

Sevin 4F.....p.114

Flea Beetles

Alias 4F.....p.20

D.Z.N Diazinon.....p.44

Sevin 4F.....p.113

Leafhoppers

Alias 4F.....p.20

Safer Insecticidal Soap.....p.103

Sevin 4F.....p.113

Leafminers

D.Z.N Diazinon.....p.44

Plant Bugs

Safer Insecticidal Soap.....p.103

Sevin 4F.....p.114

Stink Bugs

Sevin 4F.....p.114

Whiteflies

Alias 4F.....p.20

Safer Insecticidal Soap.....p.103

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Squash, Summer

Insects, Mites and Nematodes

Aphids

Alias 4F.....p.18
 Dibrom 8 Emulsive.....p.31
 D.Z.N Diazinon.....p.45
 Lannate LV.....p.74
 Malathion 5 EC.....p.80
 Malathion 8

AQUAMUL.....p.85
 Perm-Up 3.2 EC.....p.97
 Safer Insecticidal Soap.....p.104
 Vydate L.....p.123

Armyworms

Dibrom 8 Emulsive.....p.31
 Dipel DF.....p.36
 Lannate LV.....p.74

Caterpillars

Lannate LV.....p.74
 Malathion 5 EC.....p.80
 Malathion 8

AQUAMUL.....p.85
 Sevin 4F.....p.109

Cucumber Beetles

Alias 4F.....p.18
 Dibrom 8 Emulsive.....p.31
 D.Z.N Diazinon.....p.45
 Lannate LV.....p.74
 Malathion 5 EC.....p.80
 Malathion 8

AQUAMUL.....p.85
 Perm-Up 3.2 EC.....p.97
 Sevin 4F.....p.110

Cutworms

D.Z.N Diazinon.....p.45
 Lannate LV.....p.74

Perm-Up 3.2 EC.....p.97

Flea Beetles

Lannate LV.....p.74
 Sevin 4F.....p.110

Leafhoppers

Alias 4F.....p.18
 D.Z.N Diazinon.....p.45
 Perm-Up 3.2 EC.....p.97
 Sevin 4F.....p.110

Leafminers

Dibrom 8 Emulsive.....p.32
 D.Z.N Diazinon.....p.45
 Malathion 5 EC.....p.80
 Perm-Up 3.2 EC.....p.97
 Vydate L.....p.123

Loopers

Dibrom 8 Emulsive.....p.31
 Dipel DF.....p.36
 Lannate LV.....p.74
 Perm-Up 3.2 EC.....p.97

Melonworms

Dipel DF.....p.36
 D.Z.N Diazinon.....p.45
 Lannate LV.....p.74
 Perm-Up 3.2 EC.....p.97
 Sevin 4F.....p.109

Mites

Safer Insecticidal Soap.....p.104

Nematodes

Vydate L.....p.123

Plant Bugs

Perm-Up 3.2 EC.....p.97

CROP AND PEST USE INDEX

Safer Insecticidal Soap.....p.104

Squash Bugs

Perm-Up 3.2 EC.....p.97

Sevin 4F.....p.110

Spider Mites

Dibrom 8 Emulsive.....p.31

D.Z.N Diazinon.....p.45

Malathion 5 EC.....p.80

Malathion 8

AQUAMUL.....p.85

Thrips

Alias 4F.....p.18

Dibrom 8 Emulsive.....p.31

D.Z.N Diazinon.....p.45

Malathion 8

AQUAMUL.....p.85

Whiteflies

Alias 4F.....p.18

Safer Insecticidal Soap.....p.104

Diseases

Alternaria Leaf Blight

Bravo 720.....p.25

Alternaria Leaf Spot

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Angular Leaf Spot

Kocide 3000.....p.63

Anthracnose

Bravo 720.....p.24

Equis 720.....p.52

Manzate Pro Stick.....p.90

Cercospora Leaf Spot

Bravo 720.....p.25

Manzate Pro Stick.....p.90

Downy Mildew

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.92

Gummy Stem Blight

Bravo 720.....p.25

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Powdery Mildew

Bravo 720.....p.25

Kocide 3000.....p.63

Scab

Bravo 720.....p.25

Manzate Pro Stick.....p.90

Target Spot

Bravo 720.....p.24

Equis 720.....p.52

Watermelon Bacterial Fruit Blotch

Kocide 3000.....p.63

Other

CROP AND PEST USE INDEX

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Squash, Winter

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.45
Malathion 5 EC.....	p.80
Malathion 8	
AQUAMUL.....	p.86
Perm-Up 3.2 EC.....	p.97
Vydate L.....	p.123

Armyworms

Dipel DF.....	p.36
---------------	------

Caterpillars

Malathion 5 EC.....	p.80
Malathion 8	
AQUAMUL.....	p.86
Sevin 4F.....	p.109

Cucumber Beetles

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.45
Malathion 5 EC.....	p.80
Malathion 8	
AQUAMUL.....	p.86
Perm-Up 3.2 EC.....	p.97
Sevin 4F.....	p.110

Cutworms

D.Z.N Diazinon.....	p.45
Perm-Up 3.2 EC.....	p.97

Flea Beetles

Sevin 4F.....	p.110
---------------	-------

Leafhoppers

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.45
Perm-Up 3.2 EC.....	p.97
Sevin 4F.....	p.110

Leafminers

D.Z.N Diazinon.....	p.45
Malathion 5 EC.....	p.80
Perm-Up 3.2 EC.....	p.97
Vydate L.....	p.123

Loopers

Dipel DF.....	p.36
Perm-Up 3.2 EC.....	p.97

Melonworms

Dipel DF.....	p.36
D.Z.N Diazinon.....	p.45
Perm-Up 3.2 EC.....	p.97
Sevin 4F.....	p.109

Nematodes

Vydate L.....	p.123
---------------	-------

Plant Bugs

Perm-Up 3.2 EC.....	p.97
---------------------	------

Squash Bugs

Perm-Up 3.2 EC.....	p.97
Sevin 4F.....	p.110

Spider Mites

D.Z.N Diazinon.....	p.45
Malathion 5 EC.....	p.80
Malathion 8	
AQUAMUL.....	p.86

Thrips

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.45
Malathion 8	
AQUAMUL.....	p.86

Whiteflies

CROP AND PEST USE INDEX

Alias 4F.....p.18

Diseases

Alternaria Leaf Blight

Bravo 720.....p.25

Equis 720.....p.53

Alternaria Leaf Spot

Equis 720.....p.53

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Angular Leaf Spot

Kocide 3000.....p.63

Anthracnose

Bravo 720.....p.24

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Cercospora Leaf Spot

Bravo 720.....p.25

Equis 720.....p.53

Manzate Pro Stick.....p.90

Downy Mildew

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Gummy Stem Blight

Bravo 720.....p.25

Equis 720.....p.53

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Powdery Mildew

Bravo 720.....p.25

Equis 720.....p.53

Kocide 3000.....p.63

Scab

Bravo 720.....p.25

Equis 720.....p.53

Manzate Pro Stick.....p.90

Target Spot

Bravo 720.....p.24

Equis 720.....p.52

Watermelon Bacterial Fruit Blotch

Kocide 3000.....p.63

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Sweet Potato

Insects, Mites and Nematodes

Aphids

Alias 4F.....p.21

Armyworms

Sevin 4F.....p.114

Beetles

Sevin 4F.....p.114

Caterpillars

Sevin 4F.....p.114

Cucumber Beetles

Alias 4F.....p.21

Sevin 4F.....p.114

Flea Beetles

D.Z.N Diazinon.....p.45

Sevin 4F.....p.114

Leafhoppers

Alias 4F.....p.21

Malathion 5 EC.....p.80

Malathion 8

AQUAMUL.....p.86

Leafminers

Malathion 5 EC.....p.80

Malathion 8

AQUAMUL.....p.86

Nematodes

Vydate L.....p.127

Weevils

Sevin 4F.....p.114

Whiteflies

Alias 4F.....p.21

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Tangerine

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Vydate L.....	p.125

Beetles

Sevin 4F.....	p.108
Vydate L.....	p.125

Black Flies

Alias 4F.....	p.18
---------------	------

Cutworms

Dipel DF.....	p.37
Sevin 4F.....	p.107

Flea Beetles

Alias 4F.....	p.18
---------------	------

Leafhoppers

Alias 4F.....	p.18
Vydate L.....	p.125

Leafminers

Alias 4F.....	p.18
Vydate L.....	p.125

Leafrollers

Dibrom 8 Emulsive.....	p.31
Dipel DF.....	p.37
Sevin 4F.....	p.107

Mealybugs

Alias 4F.....	p.18
Vydate L.....	p.125

Mites

Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Sevin 4F.....	p.107
Vendex 50WP.....	p.122
Vydate L.....	p.124/125

Nematodes

Vydate L.....	p.125
---------------	-------

Plant Bugs

Sevin 4F.....	p.107
---------------	-------

Scale Crawlers

Dibrom 8 Emulsive.....	p.31
------------------------	------

Scales

Alias 4F.....	p.18
Dimethoate 4 EC.....	p.33
Malathion 5 EC.....	p.77
Malathion 8	
AQUAMUL.....	p.82
Sevin 4F.....	p.107
Vydate L.....	p.125

Sharpshooters

Alias 4F.....	p.18
---------------	------

Thrips

Alias 4F.....	p.18
Dibrom 8 Emulsive.....	p.31
Dimethoate 4 EC.....	p.34
Malathion 5 EC.....	p.77
Vydate L.....	p.124/125

Weevils

Sevin 4F.....	p.108
---------------	-------

Whiteflies

Alias 4F.....	p.18
Dimethoate 4 EC.....	p.34

CROP AND PEST USE INDEX

Vydate L.....p.125

Diseases

Alternaria

Kocide 3000.....p.60

Black Spot

Kocide 3000.....p.62

Brown Rot

Kocide 3000.....p.61

Brown Spot

Kocide 3000.....p.60

Citrus Canker

Kocide 3000.....p.61

Melanose

Kocide 3000.....p.60

Phytophthora

Kocide 3000.....p.61

Phytophthora Foot Rot

Kocide 3000.....p.61

Scab

Kocide 3000.....p.60

Septoria Spot

Kocide 3000.....p.61

Other

Slugs and Snails

Deadline GT.....p.48

Sluggo Snail & Slug Bait.....p.119

Taro

Consult your agricultural extension agent for information on pesticide use for this crop.

Tomato

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.19
Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.47
Lannate LV.....	p.75
Malathion 5 EC.....	p.80
Malathion 8	
AQUAMUL.....	p.86
Safer Insecticidal Soap.....	p.104
Vydate L.....	p.127

Armyworms

Dipel DF.....	p.38
D.Z.N Diazinon.....	p.47
Lannate LV.....	p.75
Malathion 5 EC.....	p.81
Malathion 8	
AQUAMUL.....	p.86
Perm-Up 3.2 EC.....	p.100
Sevin 4F.....	p.110
Sevin SL.....	p.117

Beetles

Perm-Up 3.2 EC.....	p.100
Sevin 4F.....	p.110
Vydate L.....	p.127

Caterpillars

Dipel DF.....	p.38
Lannate LV.....	p.75
Malathion 5 EC.....	p.81
Malathion 8	
AQUAMUL.....	p.86
Sevin 4F.....	p.110

Colorado Potato Beetle

Alias 4F.....	p.19
---------------	------

Cucumber Beetles

D.Z.N Diazinon.....	p.47
---------------------	------

Cutworms

Dipel DF.....	p.38
D.Z.N Diazinon.....	p.46
Lannate LV.....	p.75
Perm-Up 3.2 EC.....	p.100
Sevin 4F.....	p.111
Sevin SL.....	p.118

Flea Beetles

Sevin 4F.....	p.110
Sevin SL.....	p.117

Leafhoppers

Alias 4F.....	p.19
Dimethoate 4 EC.....	p.35
Sevin 4F.....	p.110
Sevin SL.....	p.117

Leafminers

Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.47
Perm-Up 3.2 EC.....	p.100
Vydate L.....	p.127

Loopers

Dipel DF.....	p.38
Lannate LV.....	p.75
Perm-Up 3.2 EC.....	p.100

Mites

Safer Insecticidal Soap.....	p.104
------------------------------	-------

Nematodes

Vydate L.....	p.127
---------------	-------

Plant Bugs

Sevin 4F.....	p.110
---------------	-------

Spider Mites

CROP AND PEST USE INDEX

Malathion 5 EC.....p.80
Malathion 8
AQUAMUL.....p.86

Stink Bugs

Sevin 4F.....p.110
Sevin SL.....p.117

Thrips

Sevin 4F.....p.110
Sevin SL.....p.117

Whiteflies

Alias 4F.....p.19
Safer Insecticidal Soap.....p.104

Diseases

Alternaria Fruit Rot (Black Mold)

Bravo 720.....p.28
Equis 720.....p.57

Anthracnose

Bravo 720.....p.28
Equis 720.....p.57
Kocide 3000.....p.66

Bacterial Speck

Kocide 3000.....p.66

Bacterial Spot

Kocide 3000.....p.66

Botrytis Blight (Gray Mold)

Bravo 720.....p.27/28
Equis 720.....p.57
Equis 720.....p.57
Kocide 3000.....p.66

Early Blight

Bravo 720.....p.27
Equis 720.....p.57

Fruit Rot

Equis 720.....p.57

Gray Leaf Spot

Bravo 720.....p.27
Equis 720.....p.57

Late Blight

Equis 720.....p.57
Kocide 3000.....p.66

Rhizoctonia Fruit Rot

Bravo 720.....p.28
Equis 720.....p.57

Septoria Spot

Bravo 720.....p.27
Equis 720.....p.57
Kocide 3000.....p.66

Target Spot

Bravo 720.....p.27
Equis 720.....p.57

Other

Slugs and Snails

Deadline GT.....p.48
Sluggo Snail & Slug Bait.....p.120

Watermelon

Insects, Mites and Nematodes

Aphids

Alias 4F.....	p.18
Dimethoate 4 EC.....	p.35
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.73
Malathion 5 EC.....	p.81
Malathion 8	
AQUAMUL.....	p.87
Perm-Up 3.2 EC.....	p.97
Safer Insecticidal Soap.....	p.104
Vydate L.....	p.123

Armyworms

Dipel DF.....	p.36
Lannate LV.....	p.73

Caterpillars

Lannate LV.....	p.73
-----------------	------

Cucumber Beetles

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.42
Lannate LV.....	p.73
Malathion 5 EC.....	p.81
Malathion 8	
AQUAMUL.....	p.87
Perm-Up 3.2 EC.....	p.97

Cutworms

D.Z.N Diazinon.....	p.41
Lannate LV.....	p.73

Flea Beetles

Lannate LV.....	p.73
-----------------	------

Leafhoppers

Alias 4F.....	p.18
D.Z.N Diazinon.....	p.42
Malathion 5 EC.....	p.81

Malathion 8

AQUAMUL.....p.87

Perm-Up 3.2 EC.....p.97

Leafminers

Dimethoate 4 EC.....p.35

D.Z.N Diazinon.....p.42

Malathion 5 EC.....p.81

Malathion 8

AQUAMUL.....p.87

Perm-Up 3.2 EC.....p.97

Vydate L.....p.123

Loopers

Dipel DF.....p.36

Lannate LV.....p.73

Perm-Up 3.2 EC.....p.97

Melonworms

Dipel DF.....p.36

D.Z.N Diazinon.....p.42

Lannate LV.....p.73

Perm-Up 3.2 EC.....p.97

Mites

Safer Insecticidal Soap.....p.104

Nematodes

Vydate L.....p.123

Plant Bugs

Perm-Up 3.2 EC.....p.97

Safer Insecticidal Soap.....p.104

Squash Bugs

Perm-Up 3.2 EC.....p.97

Spider Mites

D.Z.N Diazinon.....p.42

Malathion 5 EC.....p.81

CROP AND PEST USE INDEX

Malathion 8

AQUAMUL.....p.87

Thrips

Alias 4F.....p.18

D.Z.N Diazinon.....p.42

Whiteflies

Alias 4F.....p.18

Safer Insecticidal Soap.....p.104

Diseases

Alternaria Leaf Blight

Bravo 720.....p.25

Equis 720.....p.53

Alternaria Leaf Spot

Equis 720.....p.53

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Angular Leaf Spot

Kocide 3000.....p.63

Anthracnose

Bravo 720.....p.24

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Cercospora Leaf Spot

Bravo 720.....p.25

Equis 720.....p.53

Manzate Pro Stick.....p.90

Downy Mildew

Equis 720.....p.52

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Gummy Stem Blight

Bravo 720.....p.25

Equis 720.....p.53

Kocide 3000.....p.63

Manzate Pro Stick.....p.90

Powdery Mildew

Bravo 720.....p.25

Equis 720.....p.53

Kocide 3000.....p.63

Scab

Bravo 720.....p.25

Equis 720.....p.53

Manzate Pro Stick.....p.90

Target Spot

Bravo 720.....p.24

Equis 720.....p.52

Watermelon Bacterial Fruit Blotch

Kocide 3000.....p.63

Other

Slugs and Snails

Sluggo Snail & Slug Bait.....p.120

Yam

Insects, Mites and Nematodes

Aphids

Alias 4F.....p.21

Cucumber Beetles

Alias 4F.....p.21

Leafhoppers

Alias 4F.....p.21

Whiteflies

Alias 4F.....p.21

Diseases

Anthracnose

Equis 720.....p.58

6th Edition 2024

Prepared by:

Jesse Bamba, Extension Agent II, PSEP Coordinator

Elizabeth Hahn, Extension Associate I

Katherine Perez, Extension Associate I

Roland J. Quitugua, Extension Associate III

Phoebe Wall, Extension Associate III

University of Guam College of Natural & Applied Sciences Cooperative Extension & Outreach

For additional information on pesticides and pest control, please contact an agricultural extension agent at the University of Guam, Guam Cooperative Extension and Outreach, College of Natural & Applied Sciences. You may call (671) 735-2080/(671)735-2062.

Reference to a product name does not imply approval or recommendation of the product by the University of Guam, Cooperative Extension & Outreach, College of Natural & Applied Sciences, and its employees, or the U.S.D.A. to the exclusion of others which may be equally suitable.

This publication, as supported by U.S. Department of Agriculture funds through the University of Guam College of Natural & Applied Sciences, is in the public domain.

The University of Guam is a U.S. Land Grant and Sea Grant institution accredited by the WASC Senior College & University Commission. UOG is an equal opportunity provider and employer committed to diversity, equity, and inclusion through island wisdom values of *inadahi yan inagofli'e*: respect, compassion, and community.

To request this publication in alternate forms, please contact the UOG EEO/ADA/Title IX Office at (671) 735-2971/2244 or email efgoue@triton.uog.edu.

Find all UOG Extension publications at uog.edu/extension/publications.

FUNDING ACKNOWLEDGEMENT

This publication is funded by the 2019 Pesticide Safety Education Funds Management Program (PSEFMP) and 2024 Western Plant Diagnostic Network (WPDN).