WEEDS OF GUAM

Ruellia tuberosa

Weed #A2023-2 Acanthaceae






Ruellia tuberosa along roadside

Corolla tube with a darkened center



Didynamous stamens and style



Axillary inflorescence and opposite leaves



Tuberous roots

SCIENTIFIC NAME:	Ruellia tuberosa L.
COMMON NAME:	Meadow Weed, Minnieroot, Cracker Plant, Popping Pod.
SYNONYMS:	Dipteracanthus clandestinus (L.) C.Presl, Ruellia lysimachia Nees, Ruellia paniculata Scop. ex Nees, Ruellia tuberosa var. humilis M.Gómez, Ruellia picta G.Lodd, Ruellia clandestina L.
DISTRIBUTION:	Widely cultivated or naturalized in Southern USA, Central America and north of South America. Introduced to Africa, Asia and Pacific islands.

Prepared by Michael Angelo P. Fernandez, Wei Xiao, James McConnell and Mari Marutani. Factsheet: Weeds of Guam #A2023-2. November 2023. College of Natural and Applied Sciences, University of Guam Mangilao GU 96923



Young fruit with filiform calyx



Open seed capsule



Seeds



Seedlings (14-day old)



Close-up of seed

FORM:	Erect, branched herb.
ROOT:	Tuberous roots, elongate, fusiform.
STEM:	Quadrangular, deeply grooved, greenish to darkish purple, pubescent in younger parts, glabrous in older parts.
LEAVES:	Simple, blade obovate to oblanceolate, apex subacute to broadly rounded, margin undulate to approximately entire, surface glabrous or with scattered broad glossy hair along veins; oppositely arranged.
INFLORESCENCE:	3-flowered axillary dichasial cyme (or 1-flowered with two undeveloping buds), peduncle subglabrous to sparsely pubescent.
FLOWER:	Calyx : 5-lobed, filiform; corolla : mauve to bluish purple, 5-lobed, lobes orbicular to transversely elliptic, corolla diameter 4-5 cm, corolla tube narrowly funnel shaped; stamens : 4, white, included in throat, strongly didynamous, anthers not overlapping, filaments fused only at base; style : white to purple; ovary : glabrous, superior.
FRUIT:	Spindle-shaped dehiscent capsule, 2.0-2.5 cm long, brown, containing 14-34 seeds.
SEED:	Orbicular, brown, pubescent.
HABITAT:	Occur in roadsides and disturbed fields.
PROPOGATION:	Seeds, stem cuttings.
USES:	Herbal, ornamental.
MISCELLANEOUS:	Can be an invasive weed with rapid seed distribution in a habitat; well developed roots are difficult to remove.

References

Ezcurra, C. (1993). Systematics of *Ruellia* (Acanthaceae) in southern south America. Annals of the Missouri Botanical Garden, 787-845. Long, R. W. (1976). Biosystematics of *Ruellia tuberosa* L.(Acanthaceae). American Journal of Botany, 63(7), 951-959. Manzitto-Tripp, E. A., & Daniel, T. F. (2023). Phylogeny and revised classification of New World *Ruellia*. Taxon, 72(5), 1034-1056. Tripp, E. A. (2007). Evolutionary relationships within the species-rich genus *Ruellia* (Acanthaceae). Systematic Botany, 32(3), 628-649. Vollesen, K., Polhill, R. M., & Beentje, H. J. (1985). Flora of Tropical East Africa. Crown Agents for the Colonies.

Disclaimer:

Published by Agriculture and Life Sciences Division, the College of Natural & Applied Sciences, University of Guam, in cooperation with the U.S. Department of Agriculture, under Dr. Rachael T. Leon Guerrero, Director/Dean. University of Guam, CNAS, UOG Station, Mangilao, Guam 96923. The University of Guam is an equal opportunity/affirmative action institution providing programs and services to the people of Guam without regard to race, sex, gender identity and expression, age, religion, color, national origin, ancestry, disability, marital status, arrest and court record, sexual orientation, or status as a covered veteran. Reference to product name does not imply approval or recommendation of such product by the University of Guam to the exclusion of others that may be equally suitable. For more information, please contact the Dean's Office at 671-735-2001